



Full wwPDB X-ray Structure Validation Report i

Jun 13, 2024 – 12:57 AM EDT

PDB ID : 4C8S
Title : Crystal structure of the C-terminal region of yeast Ctf4
Authors : Simon, A.C.; Pellegrini, L.
Deposited on : 2013-10-01
Resolution : 3.00 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 9447 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA POLYMERASE ALPHA-BINDING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	425	Total	C 3425	N 2199	O 568	S 643	15	0	1	0
1	B	437	Total	C 3519	N 2258	O 585	S 660	16	0	1	0
1	C	296	Total	C 2405	N 1562	O 392	S 440	11	0	1	0

There are 63 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	450	MET	-	expression tag	UNP Q01454
A	451	GLY	-	expression tag	UNP Q01454
A	452	SER	-	expression tag	UNP Q01454
A	453	SER	-	expression tag	UNP Q01454
A	454	HIS	-	expression tag	UNP Q01454
A	455	HIS	-	expression tag	UNP Q01454
A	456	HIS	-	expression tag	UNP Q01454
A	457	HIS	-	expression tag	UNP Q01454
A	458	HIS	-	expression tag	UNP Q01454
A	459	HIS	-	expression tag	UNP Q01454
A	460	SER	-	expression tag	UNP Q01454
A	461	GLN	-	expression tag	UNP Q01454
A	462	ASP	-	expression tag	UNP Q01454
A	463	PRO	-	expression tag	UNP Q01454
A	464	GLU	-	expression tag	UNP Q01454
A	465	ASN	-	expression tag	UNP Q01454
A	466	LEU	-	expression tag	UNP Q01454
A	467	TYR	-	expression tag	UNP Q01454
A	468	PHE	-	expression tag	UNP Q01454
A	469	GLN	-	expression tag	UNP Q01454
A	470	GLY	-	expression tag	UNP Q01454
B	450	MET	-	expression tag	UNP Q01454
B	451	GLY	-	expression tag	UNP Q01454

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	452	SER	-	expression tag	UNP Q01454
B	453	SER	-	expression tag	UNP Q01454
B	454	HIS	-	expression tag	UNP Q01454
B	455	HIS	-	expression tag	UNP Q01454
B	456	HIS	-	expression tag	UNP Q01454
B	457	HIS	-	expression tag	UNP Q01454
B	458	HIS	-	expression tag	UNP Q01454
B	459	HIS	-	expression tag	UNP Q01454
B	460	SER	-	expression tag	UNP Q01454
B	461	GLN	-	expression tag	UNP Q01454
B	462	ASP	-	expression tag	UNP Q01454
B	463	PRO	-	expression tag	UNP Q01454
B	464	GLU	-	expression tag	UNP Q01454
B	465	ASN	-	expression tag	UNP Q01454
B	466	LEU	-	expression tag	UNP Q01454
B	467	TYR	-	expression tag	UNP Q01454
B	468	PHE	-	expression tag	UNP Q01454
B	469	GLN	-	expression tag	UNP Q01454
B	470	GLY	-	expression tag	UNP Q01454
C	450	MET	-	expression tag	UNP Q01454
C	451	GLY	-	expression tag	UNP Q01454
C	452	SER	-	expression tag	UNP Q01454
C	453	SER	-	expression tag	UNP Q01454
C	454	HIS	-	expression tag	UNP Q01454
C	455	HIS	-	expression tag	UNP Q01454
C	456	HIS	-	expression tag	UNP Q01454
C	457	HIS	-	expression tag	UNP Q01454
C	458	HIS	-	expression tag	UNP Q01454
C	459	HIS	-	expression tag	UNP Q01454
C	460	SER	-	expression tag	UNP Q01454
C	461	GLN	-	expression tag	UNP Q01454
C	462	ASP	-	expression tag	UNP Q01454
C	463	PRO	-	expression tag	UNP Q01454
C	464	GLU	-	expression tag	UNP Q01454
C	465	ASN	-	expression tag	UNP Q01454
C	466	LEU	-	expression tag	UNP Q01454
C	467	TYR	-	expression tag	UNP Q01454
C	468	PHE	-	expression tag	UNP Q01454
C	469	GLN	-	expression tag	UNP Q01454
C	470	GLY	-	expression tag	UNP Q01454


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	39	Total O 39 39	0	0
2	B	42	Total O 42 42	0	0
2	C	17	Total O 17 17	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA POLYMERASE ALPHA-BINDING PROTEIN



- Molecule 1: DNA POLYMERASE ALPHA-BINDING PROTEIN



- Molecule 1: DNA POLYMERASE ALPHA-BINDING PROTEIN



SER	LYS		
GLN	VAL		
ASP	LEU		
ARG	SER	M774	
ALA	LEU	EF75	
LEU	LEU	1776	
THR			
ALA	LEU		ARG
ALA	THR		MET
ALA	MET		PRO
VAL	ASP		VAL
	LYS		PHE
	THR		VAL
TLE	LEU		VAL
SER	GLU		LYS
GLU	ASN		SER
ARG	ASP		SER
ALA	GLY		LYS
GLU	GLU		LYS
LEU	MET		LEU
PRO	TYR		GLU
SER	GLY		GLU
LEU	ASN		ASN
ASN	ASN		ASN
VAL	GLU		LYS
	LYS		ALA
	GLU		ALA
TLE	VAL		ILE
ASN	LEU		LEU
ALA	ALA		ASN
TLE	ALA		LYS
ALA	ALA		ASN
ARG	LEU		ASN
GLU	ASN		GLU
ALA	GLY		ILE
ARG	ALA		GLY
TYR	TYR		ALA
GLU	ASP		ASP
GLN	LYS		THR
GLN	ALA		GLU
ALA	LEU		ALA
LEU	LEU		CYS
LYS	ARG		GLU
	LEU		SER
	ALA		ILE
	GLU		ASP
	GLN		GLN
	ASP		ILE
	SER		ASN
	ALA		VAL
	CYS		GLU
	GLU		SER
	ILE		LYS
	ASP		MET
	GLN		ALA
	ASP		ALA
	SER		LEU
	ALA		GLU
	LEU		LEU
	ALA		ALA
	ALA		TYR
	HIS		GLU
	TYR		LEU
	LEU		ARG

4 Data and refinement statistics i

Property	Value	Source
Space group	P 2 1 21	Depositor
Cell constants a, b, c, α , β , γ	89.06 Å 100.25 Å 219.52 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.87 – 3.00 48.87 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.87-3.00) 99.9 (48.87-3.00)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.37 (at 3.01 Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R , R_{free}	0.167 , 0.205 0.167 , 0.203	Depositor DCC
R_{free} test set	2014 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	55.1	Xtriage
Anisotropy	0.223	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.5	EDS
L-test for twinning ²	$< L > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9447	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.23	0/3510	0.40	0/4752
1	B	0.23	0/3605	0.40	0/4879
1	C	0.23	0/2481	0.39	0/3370
All	All	0.23	0/9596	0.40	0/13001

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3425	0	3369	22	0
1	B	3519	0	3477	25	0
1	C	2405	0	2326	23	0
2	A	39	0	0	0	1
2	B	42	0	0	0	1
2	C	17	0	0	0	0
All	All	9447	0	9172	65	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:510:GLN:HG2	1:C:531:LEU:HD23	1.74	0.69
1:C:728:ILE:HD11	1:C:742:VAL:HG23	1.79	0.65
1:C:722:LEU:HD21	1:C:774:MET:HE2	1.84	0.59
1:A:728:ILE:HD11	1:A:742:VAL:HG23	1.83	0.58
1:B:899:VAL:HG13	1:B:911:VAL:HG13	1.86	0.57
1:A:716:SER:O	1:C:653:ARG:NH1	2.39	0.55
1:A:533:GLY:O	1:A:548:GLN:NE2	2.40	0.54
1:A:548:GLN:HG3	1:A:551:THR:H	1.73	0.54
1:A:511:TYR:HB2	1:A:530:ASP:HB3	1.90	0.52
1:C:560:ASP:HB3	1:C:563:HIS:HD2	1.74	0.52
1:A:899:VAL:HG13	1:A:911:VAL:HG13	1.91	0.52
1:B:511:TYR:HB2	1:B:530:ASP:HB3	1.93	0.51
1:A:717:LYS:HD3	1:C:648:LYS:HB3	1.92	0.51
1:C:511:TYR:HB2	1:C:530:ASP:HB3	1.93	0.51
1:B:626:PHE:HE1	1:B:638:TYR:HB2	1.76	0.51
1:A:846:GLU:OE2	1:A:852:ASN:ND2	2.44	0.50
1:C:506:LYS:HG2	1:C:511:TYR:CE2	2.46	0.50
1:B:533:GLY:O	1:B:548:GLN:NE2	2.45	0.50
1:B:548:GLN:HG3	1:B:551:THR:H	1.76	0.49
1:B:789:GLU:HG3	1:B:818:PRO:HG3	1.94	0.49
1:A:490:ASP:HB3	1:A:506:LYS:HB2	1.93	0.49
1:A:790:ASN:HD21	1:A:815:ILE:HG23	1.77	0.49
1:C:597:PHE:HB3	1:C:610:GLU:HB3	1.93	0.49
1:A:587:ARG:CZ	1:A:607:PHE:HE2	2.26	0.49
1:B:790:ASN:HD21	1:B:815:ILE:HG23	1.78	0.49
1:A:866:LEU:HD21	1:A:888:GLU:HB2	1.96	0.48
1:C:642:GLU:HG3	1:C:651:TYR:HE2	1.78	0.48
1:B:665:ASN:H	1:B:668:MET:HG3	1.78	0.47
1:A:507:ASN:HB3	1:A:510:GLN:HG3	1.95	0.47
1:C:755:ILE:HG23	1:C:769:PRO:HG2	1.97	0.47
1:A:508:SER:OG	1:A:509:GLU:N	2.45	0.47
1:B:818:PRO:HG2	1:B:821:MET:HB3	1.96	0.46
1:B:728:ILE:HD11	1:B:742:VAL:HG23	1.97	0.46
1:C:704:LEU:HD22	1:C:752:LEU:HD13	1.96	0.46
1:C:586:VAL:O	1:C:587:ARG:NH1	2.46	0.46
1:B:633:PHE:CG	1:C:656:PRO:HG3	2.51	0.46
1:C:548:GLN:HG3	1:C:551:THR:H	1.81	0.46
1:C:685:LYS:NZ	1:C:702:ASN:OD1	2.48	0.45
1:A:484:THR:O	1:A:492:ARG:HD2	2.17	0.44
1:A:648:LYS:HB3	1:B:717:LYS:HD3	2.00	0.44
1:C:535:ASP:OD1	1:C:535:ASP:N	2.51	0.44
1:A:726:MET:O	1:A:730:LYS:HG3	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:535:ASP:OD1	1:B:535:ASP:N	2.51	0.43
1:A:586:VAL:HG23	1:A:587:ARG:HG2	2.00	0.43
1:B:866:LEU:HD23	1:B:889:LEU:HD23	2.01	0.43
1:B:668:MET:HA	1:B:671:ASP:HB2	2.00	0.43
1:C:484:THR:O	1:C:492:ARG:HD2	2.19	0.43
1:C:615:ILE:HG12	1:C:629:HIS:CE1	2.55	0.42
1:B:484:THR:O	1:B:492:ARG:HD2	2.19	0.42
1:B:640:LEU:HD12	1:B:640:LEU:HA	1.90	0.42
1:B:652:LYS:HE3	1:B:652:LYS:HB3	1.91	0.42
1:B:538:PHE:CG	1:B:582:ALA:HA	2.55	0.42
1:B:827:TYR:CD1	1:B:866:LEU:HD13	2.54	0.42
1:B:485:PRO:HB3	1:B:679:PHE:CG	2.55	0.42
1:C:617:ALA:HB3	1:C:628:VAL:HB	2.02	0.41
1:B:866:LEU:HD21	1:B:888:GLU:HB2	2.01	0.41
1:A:789:GLU:HG3	1:A:818:PRO:HG3	2.02	0.41
1:C:507:ASN:HD21	1:C:512:SER:HB2	1.85	0.41
1:C:694:ASP:HA	1:C:695:PRO:HD3	1.93	0.41
1:A:833:LEU:HB3	1:A:858:LEU:HD21	2.03	0.40
1:B:510:GLN:NE2	1:B:529:GLU:OE1	2.55	0.40
1:A:606:PRO:HB2	1:B:828:LEU:HD11	2.03	0.40
1:A:828:LEU:HD23	1:A:828:LEU:HA	1.96	0.40
1:C:746:ALA:O	1:C:752:LEU:HD12	2.22	0.40
1:B:610:GLU:OE1	1:B:648:LYS:HD3	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:2038:HOH:O	2:B:2027:HOH:O[3_554]	2.18	0.02

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	420/478 (88%)	405 (96%)	13 (3%)	2 (0%)	29 68
1	B	434/478 (91%)	418 (96%)	15 (4%)	1 (0%)	47 82
1	C	293/478 (61%)	277 (94%)	15 (5%)	1 (0%)	41 76
All	All	1147/1434 (80%)	1100 (96%)	43 (4%)	4 (0%)	41 76

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	749	TYR
1	B	749	TYR
1	C	749	TYR
1	A	508	SER

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	378/422 (90%)	378 (100%)	0	100 100
1	B	389/422 (92%)	389 (100%)	0	100 100
1	C	267/422 (63%)	267 (100%)	0	100 100
All	All	1034/1266 (82%)	1034 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	507	ASN
1	C	555	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	425/478 (88%)	-0.39	0 [100] [100]	23, 42, 88, 118	0
1	B	437/478 (91%)	-0.35	8 (1%) 68 [40]	22, 44, 97, 141	0
1	C	296/478 (61%)	-0.27	10 (3%) 45 [19]	28, 49, 108, 145	0
All	All	1158/1434 (80%)	-0.34	18 (1%) 72 [44]	22, 45, 98, 145	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	795	ASN	3.6
1	B	794	LEU	3.6
1	B	666	SER	3.4
1	C	732	SER	3.3
1	B	927	LYS	3.2
1	C	734	GLY	3.2
1	B	792	ALA	2.9
1	C	736	GLU	2.8
1	B	796	LYS	2.6
1	C	733	GLY	2.5
1	C	738	THR	2.4
1	B	793	ILE	2.4
1	C	739	ASP	2.3
1	B	667	ASP	2.3
1	C	726	MET	2.2
1	C	729	TRP	2.2
1	C	673	ASN	2.2
1	C	509	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.