



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 05:30 AM EDT

PDB ID : 6CDD
Title : Npl4 zinc finger and MPN domains (Chaetomium thermophilum)
Authors : Bodnar, N.O.; Rapoport, T.A.
Deposited on : 2018-02-08
Resolution : 2.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3676 (2.60-2.56)
Clashscore	141614	4049 (2.60-2.56)
Ramachandran outliers	138981	3979 (2.60-2.56)
Sidechain outliers	138945	3979 (2.60-2.56)
RSRZ outliers	127900	3614 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	474	<div> <div>6%</div> <div> <div></div> <div>94%</div> </div> </div>
1	B	474	<div> <div>5%</div> <div> <div></div> <div>93%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14634 atoms, of which 7119 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Npl4 zinc finger.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	459	Total	C	H	N	O	S	0	0	0
			7228	2324	3574	625	683	22			
1	B	456	Total	C	H	N	O	S	0	0	0
			7174	2308	3545	620	679	22			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	121	Total	O	0	0
			121	121		
3	B	107	Total	O	0	0
			107	107		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Npl4 zinc finger



• Molecule 1: Npl4 zinc finger



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.86Å 72.22Å 193.54Å 90.00° 96.71° 90.00°	Depositor
Resolution (Å)	96.11 – 2.58 96.11 – 2.58	Depositor EDS
% Data completeness (in resolution range)	95.2 (96.11-2.58) 83.3 (96.11-2.58)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.73 (at 2.58Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.192 , 0.229 0.203 , 0.236	Depositor DCC
R_{free} test set	2363 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	69.5	Xtriage
Anisotropy	0.478	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 68.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14634	wwPDB-VP
Average B, all atoms (Å ²)	105.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/3748	0.43	0/5095
1	B	0.25	0/3723	0.43	0/5061
All	All	0.25	0/7471	0.43	0/10156

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3654	3574	3573	9	0
1	B	3629	3545	3544	8	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	121	0	0	1	0
3	B	107	0	0	0	0
All	All	7515	7119	7117	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:435:GLU:OE1	1:B:446:ARG:NH1	2.25	0.68
1:B:145:LEU:O	1:B:197:ARG:NH2	2.35	0.60
1:B:270:ARG:NH2	1:B:510:PHE:O	2.34	0.60
1:A:270:ARG:NH2	1:A:510:PHE:O	2.33	0.60
1:A:207:GLU:OE1	1:A:207:GLU:N	2.35	0.58
1:B:207:GLU:OE1	1:B:207:GLU:N	2.36	0.57
1:A:231:HIS:ND1	1:A:232:PRO:O	2.38	0.56
1:A:146:ASP:O	1:A:205:ARG:NH1	2.44	0.51
1:A:145:LEU:O	1:A:197:ARG:NH2	2.45	0.50
1:A:546:HIS:NE2	3:A:806:HOH:O	2.35	0.49
1:B:277:ARG:NH1	1:B:520:GLU:OE2	2.47	0.48
1:B:231:HIS:ND1	1:B:232:PRO:O	2.47	0.46
1:A:588:THR:O	1:A:591:MET:HG2	2.16	0.45
1:B:533:VAL:HG12	1:B:594:GLN:HG3	1.97	0.45
1:A:193:HIS:O	1:A:197:ARG:HG2	2.19	0.42
1:A:232:PRO:HB2	1:A:236:GLU:HB2	2.02	0.42
1:B:499:ASP:OD1	1:B:500:SER:N	2.44	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	455/474 (96%)	437 (96%)	18 (4%)	0	100	100
1	B	452/474 (95%)	434 (96%)	17 (4%)	1 (0%)	47	69
All	All	907/948 (96%)	871 (96%)	35 (4%)	1 (0%)	51	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	536	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	401/414 (97%)	400 (100%)	1 (0%)	93	98
1	B	398/414 (96%)	397 (100%)	1 (0%)	92	97
All	All	799/828 (96%)	797 (100%)	2 (0%)	92	97

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	499	ASP
1	B	499	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	459/474 (96%)	0.78	28 (6%) 21 17	62, 92, 135, 163	0
1	B	456/474 (96%)	0.67	24 (5%) 26 22	68, 92, 156, 200	0
All	All	915/948 (96%)	0.73	52 (5%) 23 20	62, 92, 143, 200	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	161	ARG	4.6
1	B	236	GLU	4.1
1	A	472	TYR	4.0
1	A	593	LEU	3.7
1	B	158	ALA	3.5
1	B	241	LYS	3.4
1	B	463	GLU	3.4
1	B	249	LEU	3.2
1	B	434	ILE	3.1
1	A	559	LYS	3.0
1	A	225	ARG	2.9
1	A	476	VAL	2.9
1	A	557	LEU	2.9
1	B	161	ARG	2.8
1	A	285	ILE	2.8
1	A	359	VAL	2.8
1	B	229	SER	2.8
1	B	375	ILE	2.7
1	A	509	ILE	2.7
1	B	133	TRP	2.6
1	A	249	LEU	2.5
1	A	433	ILE	2.5
1	B	247	ILE	2.5
1	B	414	ILE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	234	TRP	2.5
1	A	238	ILE	2.5
1	B	238	ILE	2.4
1	A	547	LEU	2.4
1	B	233	GLN	2.4
1	A	589	LEU	2.3
1	B	226	ASP	2.3
1	B	489	LEU	2.3
1	A	416	ILE	2.3
1	B	205	ARG	2.2
1	A	580	LEU	2.2
1	A	477	LEU	2.2
1	B	444	LEU	2.2
1	A	135	VAL	2.1
1	A	344	VAL	2.1
1	B	405	VAL	2.1
1	A	474	ALA	2.1
1	A	530	ALA	2.1
1	B	505	PHE	2.1
1	B	433	ILE	2.1
1	B	443	MET	2.1
1	B	152	ILE	2.1
1	A	176	PRO	2.1
1	B	354	LYS	2.1
1	A	542	VAL	2.0
1	A	129	ILE	2.0
1	A	574	LEU	2.0
1	A	591	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	A	702	1/1	0.88	0.12	122,122,122,122	0
2	ZN	B	701	1/1	0.90	0.13	143,143,143,143	0
2	ZN	B	702	1/1	0.90	0.04	125,125,125,125	0
2	ZN	A	701	1/1	0.92	0.14	92,92,92,92	0

6.5 Other polymers [i](#)

There are no such residues in this entry.