



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 1, 2025 – 04:07 PM EDT

PDB ID : 9DR9 / pdb\_00009dr9  
Title : Binary product complex of DNA polymerase iota with DNA  
Authors : Frevert, Z.; Reusch, D.; Freudenthal, B.; Washington, M.T.  
Deposited on : 2024-09-25  
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.44


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7167 (2.24-2.20)
Clashscore	180529	8096 (2.24-2.20)
Ramachandran outliers	177936	8010 (2.24-2.20)
Sidechain outliers	177891	8011 (2.24-2.20)
RSRZ outliers	164620	7166 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	420	<div> <div>16%</div> <div>67%</div> <div>19%</div> <div>•</div> <div>13%</div> </div>
2	B	19	<div> <div>5%</div> <div>42%</div> <div>37%</div> <div>11%</div> <div>11%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6877 atoms, of which 3485 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase iota.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	367	Total	C	H	N	O	S	2	0	0
			5884	1825	2992	504	542	21			

- Molecule 2 is a DNA chain called DNA (5'-D(P\*AP\*AP\*GP\*GP\*GP\*TP\*CP\*CP\*TP\*AP\*GP\*GP\*AP\*CP\*CP\*CP\*T)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	17	Total	C	H	N	O	P	0	0	0
			540	165	191	66	101	17			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	132	Total	H	O	0	0
			396	264	132		
3	B	19	Total	H	O	0	0
			57	38	19		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase iota



- Molecule 2: DNA (5'-D(P\*AP\*AP\*GP\*GP\*GP\*TP\*CP\*CP\*TP\*AP\*GP\*GP\*AP\*CP\*CP\*C P\*T)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.10Å 97.10Å 202.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.76 – 2.21 43.76 – 2.21	Depositor EDS
% Data completeness (in resolution range)	97.3 (43.76-2.21) 97.2 (43.76-2.21)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.29 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.255 , 0.297 0.255 , 0.297	Depositor DCC
$R_{free}$ test set	1437 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.9	Xtriage
Anisotropy	0.279	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 32.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6877	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.66	2/2931 (0.1%)	0.97	3/3948 (0.1%)
2	B	1.26	5/391 (1.3%)	1.36	6/601 (1.0%)
All	All	0.76	7/3322 (0.2%)	1.03	9/4549 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	DA	C1'-N9	8.63	1.63	1.46
2	B	3	DA	C4'-C3'	6.03	1.64	1.52
1	A	298	PRO	CA-C	-5.85	1.48	1.52
2	B	4	DA	P-O5'	-5.64	1.49	1.60
1	A	348	ARG	CG-CD	5.37	1.68	1.52
2	B	3	DA	C5'-C4'	5.09	1.62	1.52
2	B	3	DA	N9-C4	5.03	1.48	1.38

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	DA	C4'-C3'-O3'	-12.70	90.95	110.00
2	B	3	DA	C2'-C3'-O3'	9.40	125.60	111.50
2	B	3	DA	P-O3'-C3'	9.16	133.94	120.20
2	B	3	DA	O3'-P-O5'	-6.71	93.93	104.00
2	B	3	DA	O4'-C1'-N9	-5.96	99.46	108.40
1	A	385	ASP	N-CA-CB	5.56	118.39	110.16
1	A	288	GLU	CB-CG-CD	-5.54	103.18	112.60
1	A	348	ARG	CG-CD-NE	5.21	123.47	112.00
2	B	4	DA	O5'-C5'-C4'	-5.01	103.29	110.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2892	2992	2980	79	0
2	B	349	191	191	11	0
3	A	132	264	0	2	0
3	B	19	38	0	2	0
All	All	3392	3485	3171	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:245:LYS:HE3	2:B:15:DA:H5''	1.53	0.90
1:A:62:LEU:HD13	1:A:78:LEU:HD23	1.60	0.80
1:A:64:VAL:HG13	1:A:65:THR:HG22	1.64	0.77
2:B:6:DG:H4'	3:B:101:HOH:O	1.87	0.74
1:A:325:LEU:HD21	1:A:387:LEU:HD11	1.69	0.73
1:A:325:LEU:HD21	1:A:387:LEU:CD1	2.23	0.69
1:A:69:GLU:O	1:A:73:LEU:HD13	1.94	0.67
1:A:312:SER:OG	1:A:317:ALA:HB2	1.94	0.66
1:A:69:GLU:O	1:A:73:LEU:CD1	2.43	0.66
1:A:35:LEU:HD21	1:A:187:MET:CE	2.25	0.66
1:A:27:SER:O	1:A:172:ARG:NH2	2.30	0.64
1:A:47:ASN:OD1	1:A:49:GLU:HB2	1.98	0.63
1:A:382:PRO:O	1:A:386:ILE:HD12	1.98	0.63
1:A:283:LYS:HE3	1:A:288:GLU:OE1	1.98	0.63
1:A:317:ALA:HA	3:A:512:HOH:O	1.99	0.62
1:A:310:LYS:HD3	1:A:349:TYR:HB2	1.80	0.62
1:A:267:LYS:HD3	1:A:267:LYS:H	1.67	0.60
1:A:148:LEU:O	1:A:148:LEU:HD23	2.03	0.58
1:A:229:LEU:O	1:A:232:SER:OG	2.21	0.57
1:A:308:PHE:HB2	1:A:311:CYS:HB2	1.85	0.56
1:A:35:LEU:HD21	1:A:187:MET:HE1	1.88	0.55
1:A:196:CYS:SG	1:A:214:LYS:O	2.65	0.55
2:B:4:DA:H5'	2:B:4:DA:N3	2.21	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:LYS:O	1:A:311:CYS:N	2.40	0.55
1:A:65:THR:HB	2:B:19:DT:O3'	2.06	0.54
1:A:267:LYS:HD3	1:A:267:LYS:N	2.23	0.54
1:A:77:LYS:O	1:A:78:LEU:HB2	2.07	0.54
1:A:315:VAL:CG1	1:A:315:VAL:O	2.56	0.53
1:A:275:ILE:HD12	1:A:275:ILE:H	1.73	0.53
1:A:380:MET:O	1:A:384:VAL:HG23	2.10	0.52
1:A:388:MET:HE3	1:A:388:MET:HA	1.92	0.52
1:A:35:LEU:HD21	1:A:187:MET:HE3	1.93	0.51
1:A:74:GLY:O	1:A:76:LYS:HG2	2.11	0.50
1:A:90:GLN:OE1	1:A:90:GLN:N	2.35	0.50
1:A:308:PHE:CD1	1:A:404:THR:HA	2.47	0.50
1:A:73:LEU:HD12	1:A:73:LEU:N	2.27	0.50
1:A:148:LEU:HD23	1:A:148:LEU:C	2.37	0.49
1:A:266:PRO:HD2	1:A:267:LYS:NZ	2.28	0.49
1:A:315:VAL:O	1:A:315:VAL:HG12	2.12	0.49
1:A:384:VAL:O	1:A:388:MET:HG2	2.11	0.49
1:A:379:VAL:C	1:A:382:PRO:HD2	2.39	0.48
2:B:6:DG:C4'	3:B:101:HOH:O	2.53	0.48
1:A:161:GLN:OE1	1:A:222:LEU:HB2	2.15	0.47
1:A:385:ASP:CB	3:A:523:HOH:O	2.63	0.47
1:A:36:ASP:O	1:A:37:CYS:C	2.57	0.47
1:A:66:CYS:HB2	1:A:70:ALA:HB3	1.97	0.47
1:A:309:LYS:O	1:A:310:LYS:C	2.57	0.47
2:B:3:DA:O5'	2:B:3:DA:N3	2.48	0.46
1:A:361:GLN:HG2	2:B:12:DA:OP2	2.15	0.46
2:B:8:DT:H2''	2:B:9:DC:H5'	1.96	0.46
1:A:267:LYS:H	1:A:267:LYS:CD	2.29	0.46
1:A:35:LEU:CD2	1:A:187:MET:HE1	2.45	0.45
1:A:318:LYS:O	1:A:322:GLU:HG3	2.17	0.45
1:A:325:LEU:CD2	1:A:387:LEU:CD1	2.94	0.45
2:B:15:DA:H2'	2:B:16:DC:C6	2.52	0.45
1:A:107:TYR:O	1:A:111:GLU:HG3	2.16	0.45
1:A:69:GLU:O	1:A:73:LEU:HD12	2.17	0.44
1:A:245:LYS:HD2	1:A:245:LYS:C	2.42	0.44
1:A:222:LEU:HD23	1:A:222:LEU:N	2.32	0.44
1:A:325:LEU:HB3	1:A:380:MET:HE1	1.99	0.44
1:A:255:ILE:HD11	1:A:264:PHE:CG	2.53	0.44
2:B:3:DA:N3	2:B:3:DA:C5'	2.81	0.44
2:B:8:DT:H2'	2:B:9:DC:C6	2.53	0.44
1:A:144:GLN:H	1:A:147:GLU:CD	2.26	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:312:SER:HG	1:A:317:ALA:HB2	1.81	0.43
1:A:309:LYS:C	1:A:311:CYS:N	2.76	0.43
1:A:316:GLU:O	1:A:316:GLU:OE1	2.37	0.43
1:A:275:ILE:O	1:A:279:GLN:HB2	2.19	0.43
1:A:204:LEU:HD22	1:A:284:LEU:CD2	2.48	0.42
1:A:147:GLU:O	1:A:150:ALA:N	2.44	0.42
1:A:314:GLU:HB3	1:A:316:GLU:HG3	2.01	0.42
1:A:51:LYS:O	1:A:52:ASP:HB2	2.20	0.42
1:A:248:LYS:HE3	1:A:248:LYS:HB3	1.65	0.42
1:A:307:SER:O	1:A:308:PHE:HB3	2.20	0.42
1:A:309:LYS:O	1:A:311:CYS:SG	2.76	0.42
1:A:266:PRO:HD2	1:A:267:LYS:HZ1	1.83	0.42
1:A:88:CYS:SG	1:A:91:LEU:HB2	2.60	0.42
1:A:341:THR:OG1	1:A:412:ASN:HB2	2.20	0.41
1:A:253:LEU:HD11	1:A:272:GLU:HG2	2.01	0.41
1:A:61:TYR:CD2	1:A:62:LEU:HD23	2.56	0.41
1:A:132:LEU:HD21	1:A:183:MET:HE1	2.03	0.41
1:A:144:GLN:O	1:A:145:SER:C	2.61	0.41
1:A:204:LEU:HD22	1:A:284:LEU:HD22	2.03	0.41
1:A:317:ALA:HB1	1:A:406:LEU:HD11	2.03	0.41
1:A:381:THR:N	1:A:382:PRO:CD	2.85	0.40
1:A:385:ASP:O	1:A:386:ILE:C	2.63	0.40
1:A:391:PHE:CD1	1:A:391:PHE:C	2.98	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	359/420 (86%)	338 (94%)	19 (5%)	2 (1%)	22	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	310	LYS
1	A	37	CYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	334/376 (89%)	329 (98%)	5 (2%)	60 73

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	LEU
1	A	65	THR
1	A	73	LEU
1	A	149	SER
1	A	318	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	ASN
1	A	256	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	367/420 (87%)	1.01	67 (18%) <b>4</b> <b>4</b>	31, 54, 92, 107	0
2	B	17/19 (89%)	0.41	1 (5%) 29 27	36, 44, 59, 88	0
All	All	384/439 (87%)	0.99	68 (17%) <b>4</b> <b>4</b>	31, 53, 91, 107	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	315	VAL	4.4
2	B	3	DA	4.4
1	A	317	ALA	4.3
1	A	244	TYR	4.3
1	A	68	TYR	4.2
1	A	350	SER	4.2
1	A	316	GLU	4.2
1	A	349	TYR	4.1
1	A	370	GLN	4.0
1	A	77	LYS	4.0
1	A	311	CYS	4.0
1	A	367	HIS	4.0
1	A	385	ASP	3.9
1	A	214	LYS	3.9
1	A	275	ILE	3.9
1	A	215	PRO	3.9
1	A	314	GLU	3.9
1	A	336	GLY	3.9
1	A	309	LYS	3.7
1	A	86	GLU	3.7
1	A	348	ARG	3.6
1	A	73	LEU	3.5
1	A	379	VAL	3.5
1	A	141	GLN	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	368	VAL	3.4
1	A	46	SER	3.3
1	A	394	MET	3.3
1	A	87	LYS	3.1
1	A	414	LYS	3.1
1	A	356	GLY	3.0
1	A	310	LYS	3.0
1	A	78	LEU	3.0
1	A	134	GLU	3.0
1	A	76	LYS	2.9
1	A	245	LYS	2.8
1	A	276	SER	2.8
1	A	272	GLU	2.8
1	A	369	ILE	2.8
1	A	280	ARG	2.8
1	A	236	ILE	2.7
1	A	71	ARG	2.7
1	A	148	LEU	2.6
1	A	344	LEU	2.6
1	A	332	VAL	2.6
1	A	269	LEU	2.4
1	A	312	SER	2.4
1	A	386	ILE	2.4
1	A	52	ASP	2.4
1	A	64	VAL	2.4
1	A	75	VAL	2.4
1	A	145	SER	2.4
1	A	366	SER	2.3
1	A	273	LEU	2.3
1	A	82	ARG	2.3
1	A	335	ASP	2.3
1	A	80	ASN	2.3
1	A	302	PHE	2.2
1	A	391	PHE	2.2
1	A	334	GLN	2.2
1	A	79	MET	2.2
1	A	357	ARG	2.2
1	A	146	ASP	2.2
1	A	138	LYS	2.1
1	A	364	ILE	2.1
1	A	51	LYS	2.1
1	A	49	GLU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	308	PHE	2.1
1	A	74	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.