



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 22, 2024 – 07:24 AM EDT

PDB ID : 4FET
Title : Catalytic domain of germination-specific lytic tansglycosylase SleB from *Bacillus anthracis*
Authors : Jing, X.; Heffron, J.; Popham, D.L.; Schubot, F.D.
Deposited on : 2012-05-30
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	222	<div> <div>%</div> <div> <div></div> <div>48%</div> <div>7%</div> <div>45%</div> </div> </div>
1	B	222	<div> <div>2%</div> <div> <div></div> <div>49%</div> <div>5%</div> <div>45%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2146 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spore cortex-lytic enzyme prepeptide.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	B	123	Total	C	N	O	S	Se	0	0	0
			952	610	162	177	1	2			
1	A	122	Total	C	N	O	S	Se	0	0	0
			944	606	160	175	1	2			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	32	GLY	-	expression tag	UNP Q81PQ3
B	143	MSE	LEU	engineered mutation	UNP Q81PQ3
A	32	GLY	-	expression tag	UNP Q81PQ3
A	143	MSE	LEU	engineered mutation	UNP Q81PQ3

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Na	0	0
			1	1		
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	116	Total	O	0	0
			116	116		
3	A	132	Total	O	0	0
			132	132		

i

- Molecule 1: Spore cortex-lytic enzyme prepeptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	53.85Å 64.45Å 84.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.22 – 1.91 32.22 – 1.91	Depositor EDS
% Data completeness (in resolution range)	99.8 (32.22-1.91) 99.8 (32.22-1.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.06 (at 1.91Å)	Xtriage
Refinement program	CCP4, REFMAC 5.6.0117	Depositor
R, R_{free}	0.203 , 0.249 0.205 , 0.253	Depositor DCC
R_{free} test set	1212 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å ²)	13.5	Xtriage
Anisotropy	0.066	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.55$, $\langle L^2 \rangle = 0.40$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2146	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 57.64 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.2967e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.12	3/967 (0.3%)	1.01	0/1317
1	B	1.24	3/975 (0.3%)	1.07	4/1328 (0.3%)
All	All	1.18	6/1942 (0.3%)	1.04	4/2645 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	238	TRP	CD2-CE2	6.82	1.49	1.41
1	A	238	TRP	CD2-CE2	5.64	1.48	1.41
1	A	236	TRP	CD2-CE2	5.46	1.47	1.41
1	A	217	TRP	CD2-CE2	5.39	1.47	1.41
1	B	236	TRP	CD2-CE2	5.37	1.47	1.41
1	B	204	GLU	CD-OE1	5.21	1.31	1.25

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	187	PRO	C-N-CA	-6.74	104.85	121.70
1	B	188	ARG	CG-CD-NE	-6.05	99.09	111.80
1	B	158	LEU	CB-CG-CD2	5.91	121.04	111.00
1	B	188	ARG	NE-CZ-NH2	-5.30	117.65	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	188	ARG	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	944	0	929	17	0
1	B	952	0	935	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	132	0	0	12	0
3	B	116	0	0	2	0
All	All	2146	0	1864	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:220:THR:HG22	3:A:518:HOH:O	1.73	0.88
1:A:207:LYS:CE	3:A:471:HOH:O	2.24	0.86
1:A:172:THR:HG22	3:A:469:HOH:O	1.84	0.77
1:B:205:THR:HG22	3:B:445:HOH:O	1.86	0.75
1:B:131:ASN:HB2	1:B:138:GLN:HE22	1.58	0.69
1:A:203:ASN:HD22	1:A:206:ALA:H	1.41	0.69
1:A:246:ILE:HD12	3:A:518:HOH:O	1.92	0.68
1:A:251:PHE:CD2	3:A:518:HOH:O	2.50	0.62
1:A:204:GLU:HB3	3:A:428:HOH:O	1.99	0.61
1:B:207:LYS:HZ2	1:B:207:LYS:CB	2.12	0.61
1:A:207:LYS:HZ2	1:A:207:LYS:CB	2.14	0.60
1:A:149:TYR:CZ	1:A:153:ARG:HD2	2.40	0.57
1:B:131:ASN:HB2	1:B:138:GLN:NE2	2.21	0.56
1:A:172:THR:HG21	3:A:430:HOH:O	2.09	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:158:LEU:HD23	1:B:217:TRP:CZ2	2.45	0.52
1:A:251:PHE:CG	3:A:518:HOH:O	2.67	0.48
1:B:205:THR:HG21	3:A:409:HOH:O	2.14	0.46
1:A:207:LYS:HZ2	1:A:207:LYS:HB2	1.81	0.46
1:B:221:GLY:HA2	3:B:442:HOH:O	2.16	0.45
1:A:207:LYS:NZ	3:A:471:HOH:O	2.45	0.45
1:A:207:LYS:HE2	3:A:471:HOH:O	2.04	0.43
1:A:134:ASN:HB2	3:A:494:HOH:O	2.18	0.43
1:A:207:LYS:HZ3	1:A:207:LYS:HG2	1.30	0.42
1:A:162:ALA:HB1	1:A:220:THR:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/222 (54%)	118 (98%)	2 (2%)	0	100	100
1	B	121/222 (54%)	120 (99%)	1 (1%)	0	100	100
All	All	241/444 (54%)	238 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	99/174 (57%)	97 (98%)	2 (2%)	50	47
1	B	100/174 (58%)	95 (95%)	5 (5%)	20	13
All	All	199/348 (57%)	192 (96%)	7 (4%)	31	24

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	158	LEU
1	B	188	ARG
1	B	201	THR
1	B	207	LYS
1	B	235	LYS
1	A	201	THR
1	A	207	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	138	GLN
1	A	203	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	120/222 (54%)	-0.10	2 (1%) 69 71	8, 14, 35, 42	0
1	B	121/222 (54%)	-0.19	4 (3%) 49 51	7, 12, 30, 52	0
All	All	241/444 (54%)	-0.15	6 (2%) 58 60	7, 13, 33, 52	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	236	TRP	4.7
1	B	188	ARG	3.6
1	A	236	TRP	3.0
1	A	132	VAL	2.9
1	B	131	ASN	2.8
1	B	235	LYS	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NA	B	301	1/1	0.98	0.03	10,10,10,10	0
2	NA	A	301	1/1	0.98	0.04	9,9,9,9	0

6.5 Other polymers [i](#)

There are no such residues in this entry.