



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 19, 2025 – 06:04 PM EDT

PDB ID : 4H27  
Title : Modulating the function of human serine racemase and human serine dehydratase by protein engineering  
Authors : Wang, C.Y.; Wang, A.H.  
Deposited on : 2012-09-12  
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.4


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1387 (1.30-1.30)
Clashscore	180529	1497 (1.30-1.30)
Ramachandran outliers	177936	1455 (1.30-1.30)
Sidechain outliers	177891	1455 (1.30-1.30)
RSRZ outliers	164620	1384 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2846 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-serine dehydratase/L-threonine deaminase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	318	2369	1518	401	437	1	12	0	0	0

There are 38 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-35	MET	-	expression tag	UNP P20132
A	-34	GLY	-	expression tag	UNP P20132
A	-33	SER	-	expression tag	UNP P20132
A	-32	SER	-	expression tag	UNP P20132
A	-31	HIS	-	expression tag	UNP P20132
A	-30	HIS	-	expression tag	UNP P20132
A	-29	HIS	-	expression tag	UNP P20132
A	-28	HIS	-	expression tag	UNP P20132
A	-27	HIS	-	expression tag	UNP P20132
A	-26	HIS	-	expression tag	UNP P20132
A	-25	SER	-	expression tag	UNP P20132
A	-24	SER	-	expression tag	UNP P20132
A	-23	GLY	-	expression tag	UNP P20132
A	-22	LEU	-	expression tag	UNP P20132
A	-21	VAL	-	expression tag	UNP P20132
A	-20	PRO	-	expression tag	UNP P20132
A	-19	ARG	-	expression tag	UNP P20132
A	-18	GLY	-	expression tag	UNP P20132
A	-17	SER	-	expression tag	UNP P20132
A	-16	HIS	-	expression tag	UNP P20132
A	-15	MET	-	expression tag	UNP P20132
A	-14	ALA	-	expression tag	UNP P20132
A	-13	SER	-	expression tag	UNP P20132
A	-12	MET	-	expression tag	UNP P20132
A	-11	THR	-	expression tag	UNP P20132
A	-10	GLY	-	expression tag	UNP P20132
A	-9	GLY	-	expression tag	UNP P20132

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	GLN	-	expression tag	UNP P20132
A	-7	GLN	-	expression tag	UNP P20132
A	-6	MET	-	expression tag	UNP P20132
A	-5	GLY	-	expression tag	UNP P20132
A	-4	ARG	-	expression tag	UNP P20132
A	-3	GLY	-	expression tag	UNP P20132
A	-2	SER	-	expression tag	UNP P20132
A	-1	GLU	-	expression tag	UNP P20132
A	0	PHE	-	expression tag	UNP P20132
A	65	SER	ALA	engineered mutation	UNP P20132
A	89	GLY	SER	SEE REMARK 999	UNP P20132

- Molecule 2 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	462	Total O 462 462	0	0



- Molecule 1: L-serine dehydratase/L-threonine deaminase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	158.50Å 158.50Å 59.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.30 30.00 – 1.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-1.30) 99.7 (30.00-1.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.30 (at 1.30Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.144 , 0.168 0.141 , 0.166	Depositor DCC
$R_{free}$ test set	4608 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.1	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 49.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2846	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/2392	0.74	3/3247 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	51	ARG	NE-CZ-NH2	-8.29	116.16	120.30
1	A	51	ARG	NE-CZ-NH1	7.37	123.99	120.30
1	A	51	ARG	CG-CD-NE	-5.88	99.46	111.80

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2369	0	2425	20	0
2	A	15	0	0	1	0
3	A	462	0	0	12	2
All	All	2846	0	2425	21	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:GLY:HA3	3:A:909:HOH:O	1.71	0.91
2:A:403:SO4:O2	3:A:827:HOH:O	1.93	0.85
1:A:237:GLN:HG3	3:A:943:HOH:O	1.85	0.76
1:A:67:ASN:HD21	1:A:306:SER:H	1.38	0.72
1:A:59:HIS:HE1	1:A:84:THR:OG1	1.84	0.59
1:A:293:THR:HG23	3:A:908:HOH:O	2.05	0.56
1:A:237:GLN:CG	3:A:943:HOH:O	2.49	0.55
1:A:188:VAL:O	1:A:239:HIS:HE1	1.90	0.54
1:A:59:HIS:HD2	3:A:748:HOH:O	1.90	0.53
1:A:51:ARG:NH2	3:A:788:HOH:O	2.27	0.52
1:A:307:ASN:HA	3:A:720:HOH:O	2.08	0.52
1:A:100:LYS:NZ	3:A:873:HOH:O	2.17	0.50
1:A:239:HIS:HD2	3:A:603:HOH:O	1.94	0.50
1:A:89:GLY:CA	3:A:909:HOH:O	2.43	0.47
1:A:239:HIS:CG	1:A:240:PRO:HD2	2.53	0.43
1:A:59:HIS:CE1	1:A:84:THR:OG1	2.70	0.43
1:A:188:VAL:O	1:A:239:HIS:CE1	2.71	0.42
1:A:96:ILE:HG22	1:A:100:LYS:HD2	2.02	0.41
1:A:229:GLY:HA3	3:A:607:HOH:O	2.21	0.41
1:A:70:MET:HE3	1:A:70:MET:HA	2.03	0.41
1:A:139:PRO:HA	1:A:142:TRP:CE3	2.57	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:820:HOH:O	3:A:820:HOH:O[6_556]	1.50	0.70
3:A:669:HOH:O	3:A:816:HOH:O[15_456]	2.17	0.03

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/364 (86%)	308 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/283 (87%)	245 (100%)	1 (0%)	89	74

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	HIS
1	A	67	ASN
1	A	239	HIS
1	A	290	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	LLP	A	41	1	23,24,25	1.70	6 (26%)	25,32,34	1.19	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	41	1	-	5/16/17/19	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	41	LLP	O3-C3	-4.84	1.25	1.36
1	A	41	LLP	C2-N1	2.95	1.39	1.33
1	A	41	LLP	CE-NZ	2.63	1.52	1.46
1	A	41	LLP	C4'-NZ	2.46	1.35	1.27
1	A	41	LLP	C4-C4'	2.40	1.51	1.46
1	A	41	LLP	C6-N1	2.38	1.39	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	41	LLP	C2'-C2-C3	2.36	123.56	120.80
1	A	41	LLP	C4-C4'-NZ	-2.22	113.78	124.04
1	A	41	LLP	OP4-C5'-C5	2.12	113.34	109.36
1	A	41	LLP	C5-C6-N1	-2.03	120.52	123.83

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	41	LLP	C4-C4'-NZ-CE
1	A	41	LLP	C-CA-CB-CG
1	A	41	LLP	CA-CB-CG-CD
1	A	41	LLP	CD-CE-NZ-C4'
1	A	41	LLP	C3-C4-C4'-NZ

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	401	-	4,4,4	0.28	0	6,6,6	0.47	0
2	SO4	A	402	-	4,4,4	0.41	0	6,6,6	0.79	0
2	SO4	A	403	-	4,4,4	0.43	0	6,6,6	0.99	1 (16%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	403	SO4	O4-S-O3	2.04	119.79	108.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	403	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	317/364 (87%)	-0.01	13 (4%)	42 45	5, 14, 31, 38	2 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	113	LEU	6.1
1	A	94	LEU	5.5
1	A	157	TRP	5.3
1	A	112	LEU	5.1
1	A	301	ILE	3.8
1	A	111	GLU	3.6
1	A	93	ALA	3.2
1	A	110	GLY	2.8
1	A	88	PRO	2.4
1	A	5	GLU	2.2
1	A	318	GLU	2.1
1	A	90	THR	2.1
1	A	154	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	LLP	A	41	24/25	0.98	0.07	8,15,20,22	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	403	5/5	0.83	0.15	22,27,32,33	0
2	SO4	A	402	5/5	0.85	0.15	27,31,32,32	0
2	SO4	A	401	5/5	0.97	0.07	18,20,25,27	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.