



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 29, 2025 – 09:23 AM EDT

PDB ID : 4HI3 / pdb\_00004hi3  
Title : Crystal structure of dimeric R298A mutant of SARS coronavirus main protease  
Authors : Wu, C.-G.; Chou, C.-Y.  
Deposited on : 2012-10-11  
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.43.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	 78% 16% • 5%
1	B	315	 76% 18% • •

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4912 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3C-like proteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	300	Total	C	N	O	S	0	0	0
			2320	1468	395	435	22			
1	B	301	Total	C	N	O	S	0	0	0
			2326	1471	396	437	22			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P0C6U8
A	298	ALA	ARG	engineered mutation	UNP P0C6U8
A	307	LEU	-	expression tag	UNP P0C6U8
A	308	GLU	-	expression tag	UNP P0C6U8
A	309	HIS	-	expression tag	UNP P0C6U8
A	310	HIS	-	expression tag	UNP P0C6U8
A	311	HIS	-	expression tag	UNP P0C6U8
A	312	HIS	-	expression tag	UNP P0C6U8
A	313	HIS	-	expression tag	UNP P0C6U8
A	314	HIS	-	expression tag	UNP P0C6U8
B	0	MET	-	initiating methionine	UNP P0C6U8
B	298	ALA	ARG	engineered mutation	UNP P0C6U8
B	307	LEU	-	expression tag	UNP P0C6U8
B	308	GLU	-	expression tag	UNP P0C6U8
B	309	HIS	-	expression tag	UNP P0C6U8
B	310	HIS	-	expression tag	UNP P0C6U8
B	311	HIS	-	expression tag	UNP P0C6U8
B	312	HIS	-	expression tag	UNP P0C6U8
B	313	HIS	-	expression tag	UNP P0C6U8
B	314	HIS	-	expression tag	UNP P0C6U8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	134	Total 134	O 134	0	0
2	B	132	Total 132	O 132	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: 3C-like proteinase



#### • Molecule 1: 3C-like proteinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.04Å 59.44Å 59.82Å 71.29° 73.40° 72.31°	Depositor
Resolution (Å)	30.00 – 2.09 30.00 – 2.09	Depositor EDS
% Data completeness (in resolution range)	94.7 (30.00-2.09) 94.9 (30.00-2.09)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.52 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.190 , 0.241 0.189 , 0.241	Depositor DCC
$R_{free}$ test set	1874 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.7	Xtriage
Anisotropy	0.309	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 28.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.016 for -h,-l,-k	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4912	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.73	0/2372	0.85	3/3224 (0.1%)
1	B	0.75	0/2378	0.89	3/3232 (0.1%)
All	All	0.74	0/4750	0.87	6/6456 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	154	TYR	CB-CA-C	-7.01	107.69	117.23
1	B	154	TYR	CB-CA-C	-6.40	108.52	117.23
1	A	275	GLY	N-CA-C	-6.11	105.81	112.04
1	A	165	MET	N-CA-C	5.83	117.71	108.79
1	B	148	VAL	N-CA-C	5.58	117.44	108.85
1	B	2	GLY	N-CA-C	-5.15	106.52	112.29

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2320	0	2273	39	0
1	B	2326	0	2278	47	0
2	A	134	0	0	16	0
2	B	132	0	0	13	0
All	All	4912	0	4551	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:219:PHE:HZ	2:B:509:HOH:O	1.38	1.05
1:B:212:VAL:HG23	2:B:509:HOH:O	1.58	1.02
1:A:6:MET:HB3	2:B:479:HOH:O	1.62	0.99
1:B:225:THR:HB	2:B:442:HOH:O	1.66	0.95
1:A:300:CYS:HB3	2:A:505:HOH:O	1.69	0.91
1:B:154:TYR:HB2	2:B:420:HOH:O	1.76	0.85
1:A:124:GLY:HA3	1:B:6:MET:HE3	1.61	0.83
1:B:219:PHE:CZ	2:B:509:HOH:O	2.20	0.75
1:B:109:GLY:HA2	1:B:200:ILE:HD13	1.68	0.75
1:A:207:TRP:HE3	2:A:512:HOH:O	1.71	0.73
1:A:207:TRP:CE3	2:A:512:HOH:O	2.42	0.73
1:B:186:VAL:H	1:B:192:GLN:HE22	1.37	0.72
1:A:137:LYS:O	1:B:4:ARG:NH1	2.25	0.70
1:B:66:PHE:HB2	1:B:77:VAL:HG21	1.74	0.70
1:A:4:ARG:H	1:A:299:GLN:HE22	1.41	0.69
1:B:4:ARG:H	1:B:299:GLN:HE22	1.43	0.67
1:A:145:CYS:SG	2:A:506:HOH:O	2.53	0.67
1:B:69:GLN:HB3	2:B:459:HOH:O	1.97	0.63
1:A:243:THR:HG22	1:A:246:HIS:H	1.65	0.62
1:A:140:PHE:HA	2:A:510:HOH:O	2.00	0.61
1:B:225:THR:HG22	1:B:226:THR:O	2.04	0.56
1:A:270:GLU:HG3	1:A:274:ASN:HD21	1.71	0.54
1:B:131:ARG:CD	1:B:197:ASP:OD1	2.55	0.54
1:B:23:GLY:HA3	2:B:422:HOH:O	2.07	0.53
1:A:126:TYR:CD1	1:B:4:ARG:HG2	2.44	0.53
1:A:180:LYS:HD3	2:A:525:HOH:O	2.09	0.52
1:A:167:LEU:HB3	1:A:168:PRO:CD	2.40	0.52
1:A:30:LEU:HG	1:A:148:VAL:HG11	1.91	0.51
2:A:479:HOH:O	1:B:138:GLY:HA2	2.11	0.51
1:B:131:ARG:HD2	1:B:197:ASP:OD1	2.11	0.51
1:A:6:MET:HE2	1:B:139:SER:HB3	1.93	0.50
1:A:4:ARG:HD2	1:B:126:TYR:CD1	2.47	0.49
1:B:90:LYS:HD2	2:B:429:HOH:O	2.12	0.49
1:A:3:PHE:HZ	1:A:300:CYS:SG	2.36	0.49
1:B:131:ARG:HH22	1:B:289:ASP:CG	2.21	0.49
1:A:208:LEU:HD23	2:A:512:HOH:O	2.13	0.48
1:A:276:MET:HE1	1:A:286:ILE:HA	1.95	0.48

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:4:ARG:N	1:B:299:GLN:HE22	2.11	0.48
1:B:131:ARG:HD3	1:B:197:ASP:OD1	2.14	0.48
1:A:10:SER:O	1:A:14:GLU:HG3	2.13	0.48
1:A:3:PHE:CZ	1:A:300:CYS:SG	3.07	0.48
1:B:4:ARG:NH2	2:B:528:HOH:O	2.47	0.47
1:A:132:PRO:HG2	1:A:198:THR:O	2.14	0.47
1:A:165:MET:HE3	1:A:165:MET:HB3	1.85	0.47
1:B:213:ILE:HG21	1:B:300:CYS:HB3	1.96	0.47
1:A:279:ARG:HB3	2:A:432:HOH:O	2.15	0.47
1:B:131:ARG:NH2	1:B:289:ASP:OD2	2.37	0.46
1:A:55:GLU:HG2	2:A:419:HOH:O	2.15	0.46
2:A:479:HOH:O	1:B:126:TYR:HE1	1.98	0.46
1:A:227:LEU:HD11	1:A:242:LEU:HB3	1.97	0.46
1:B:140:PHE:HD1	1:B:172:HIS:CG	2.35	0.45
1:B:125:VAL:N	2:B:479:HOH:O	2.49	0.45
1:B:176:ASP:HB2	2:B:529:HOH:O	2.16	0.45
1:B:198:THR:HG22	1:B:238:ASN:OD1	2.17	0.45
1:A:8:PHE:HB3	1:A:152:ILE:HD12	1.99	0.45
1:B:106:ILE:HD13	1:B:130:MET:HB2	1.98	0.45
1:A:218:TRP:CD1	2:A:432:HOH:O	2.70	0.44
1:B:95:ASN:HB3	1:B:98:THR:OG1	2.17	0.44
1:A:190:THR:N	2:A:446:HOH:O	2.50	0.44
1:B:106:ILE:HD11	1:B:130:MET:HE3	1.99	0.44
1:B:108:PRO:HB3	1:B:132:PRO:HA	2.00	0.44
1:A:211:ALA:HA	1:A:282:LEU:HD13	1.99	0.44
1:B:243:THR:H	1:B:246:HIS:CD2	2.36	0.44
2:A:479:HOH:O	1:B:128:CYS:HB3	2.18	0.44
1:A:276:MET:O	1:A:279:ARG:HD3	2.18	0.43
1:B:225:THR:HG22	1:B:226:THR:N	2.33	0.43
1:B:163:HIS:CE1	1:B:172:HIS:HB3	2.53	0.43
1:A:61:LYS:HA	1:A:61:LYS:HD3	1.84	0.43
1:B:40:ARG:O	1:B:43:ILE:HG12	2.19	0.43
1:B:80:HIS:HB3	1:B:89:LEU:HD23	2.01	0.42
1:B:140:PHE:HD1	1:B:172:HIS:CD2	2.36	0.42
1:B:211:ALA:HB2	1:B:282:LEU:HD22	2.00	0.42
1:A:185:PHE:HA	1:A:192:GLN:OE1	2.20	0.42
1:A:113:SER:O	1:A:149:GLY:HA2	2.20	0.42
1:B:218:TRP:CZ2	1:B:279:ARG:HB2	2.54	0.42
1:A:118:TYR:CE1	1:A:144:SER:HB3	2.55	0.42
1:B:106:ILE:H	1:B:106:ILE:HG13	1.71	0.41
1:B:265:CYS:HB2	2:B:442:HOH:O	2.21	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:270:GLU:HG3	1:A:274:ASN:ND2	2.35	0.41
1:A:281:ILE:HG12	2:A:514:HOH:O	2.20	0.40
1:A:61:LYS:HE2	2:A:435:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	298/315 (95%)	285 (96%)	13 (4%)	0	100	100
1	B	299/315 (95%)	292 (98%)	7 (2%)	0	100	100
All	All	597/630 (95%)	577 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/271 (95%)	244 (95%)	13 (5%)	20	19
1	B	258/271 (95%)	245 (95%)	13 (5%)	20	20
All	All	515/542 (95%)	489 (95%)	26 (5%)	20	20

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	LEU
1	A	67	LEU
1	A	87	LEU
1	A	94	SER
1	A	121	SER
1	A	127	GLN
1	A	165	MET
1	A	207	TRP
1	A	243	THR
1	A	268	LEU
1	A	273	GLN
1	A	279	ARG
1	A	282	LEU
1	B	4	ARG
1	B	62	SER
1	B	65	SER
1	B	75	LEU
1	B	87	LEU
1	B	106	ILE
1	B	127	GLN
1	B	131	ARG
1	B	155	ASP
1	B	178	GLU
1	B	217	ARG
1	B	268	LEU
1	B	282	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	246	HIS
1	A	273	GLN
1	A	274	ASN
1	A	299	GLN
1	B	41	HIS
1	B	69	GLN
1	B	74	GLN
1	B	127	GLN
1	B	192	GLN
1	B	246	HIS
1	B	274	ASN
1	B	299	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	300/315 (95%)	-0.19	2 (0%) 84 85	28, 43, 74, 89	0
1	B	301/315 (95%)	-0.23	3 (0%) 79 80	26, 42, 66, 86	0
All	All	601/630 (95%)	-0.21	5 (0%) 82 83	26, 42, 71, 89	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	155	ASP	2.8
1	A	154	TYR	2.8
1	B	49	MET	2.3
1	B	120	GLY	2.3
1	A	223	PHE	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.