



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 29, 2025 – 07:56 AM EDT

PDB ID : 2IA2 / pdb_00002ia2
Title : The crystal structure of a putative transcriptional regulator RHA06195 from Rhodococcus sp. RHA1
Authors : Cymborowski, M.T.; Chruszcz, M.; Skarina, T.; Onopriyenko, O.; Savchenko, A.; Edwards, A.; Joachimiak, A.; Minor, W.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2006-09-06
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 9%, green 85%, grey 6%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 85% 9% 6% </div> </div>
1	B	265	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 11%, orange 1%, yellow 7%, green 85%, grey 7%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 11% 85% 7% 7% </div> </div>
1	C	265	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 9%, orange 1%, yellow 9%, green 84%, grey 7%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 9% 84% 9% 7% </div> </div>
1	D	265	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 14%, orange 1%, yellow 9%, green 82%, grey 8%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 14% 82% 9% 8% </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative Transcriptional Regulator.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	Se	0	0	0
			1900	1194	337	364	2	3			
1	B	246	Total	C	N	O	S	Se	0	1	0
			1831	1155	319	352	2	3			
1	C	247	Total	C	N	O	S	Se	0	0	0
			1800	1129	317	349	2	3			
1	D	245	Total	C	N	O	S	Se	0	0	0
			1792	1126	315	346	2	3			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP Q2F1F8
A	133	MSE	MET	modified residue	UNP Q2F1F8
A	150	MSE	MET	modified residue	UNP Q2F1F8
A	212	MSE	MET	modified residue	UNP Q2F1F8
A	218	GLY	-	cloning artifact	UNP Q2F1F8
B	1	MSE	MET	modified residue	UNP Q2F1F8
B	133	MSE	MET	modified residue	UNP Q2F1F8
B	150	MSE	MET	modified residue	UNP Q2F1F8
B	212	MSE	MET	modified residue	UNP Q2F1F8
B	218	GLY	-	cloning artifact	UNP Q2F1F8
C	1	MSE	MET	modified residue	UNP Q2F1F8
C	133	MSE	MET	modified residue	UNP Q2F1F8
C	150	MSE	MET	modified residue	UNP Q2F1F8
C	212	MSE	MET	modified residue	UNP Q2F1F8
C	218	GLY	-	cloning artifact	UNP Q2F1F8
D	1	MSE	MET	modified residue	UNP Q2F1F8
D	133	MSE	MET	modified residue	UNP Q2F1F8
D	150	MSE	MET	modified residue	UNP Q2F1F8
D	212	MSE	MET	modified residue	UNP Q2F1F8
D	218	GLY	-	cloning artifact	UNP Q2F1F8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	214	Total 214	O 214	0	0
2	B	133	Total 133	O 133	0	0
2	C	143	Total 143	O 143	0	0
2	D	251	Total 251	O 251	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Putative Transcriptional Regulator




• Molecule 1: Putative Transcriptional Regulator




• Molecule 1: Putative Transcriptional Regulator



• Molecule 1: Putative Transcriptional Regulator





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.89Å 121.53Å 174.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.74 – 2.10 36.74 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.8 (36.74-2.10) 96.8 (36.74-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.69 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.186 , 0.223 0.201 , 0.192	Depositor DCC
R_{free} test set	3711 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtriage
Anisotropy	0.404	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8064	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.15	1/1923 (0.1%)	1.02	0/2614
1	B	1.00	2/1854 (0.1%)	1.05	2/2519 (0.1%)
1	C	1.12	3/1822 (0.2%)	1.03	1/2475 (0.0%)
1	D	1.32	9/1813 (0.5%)	1.06	0/2459
All	All	1.15	15/7412 (0.2%)	1.04	3/10067 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	246	ILE	CA-CB	7.59	1.59	1.53
1	D	246	ILE	CA-C	7.33	1.58	1.52
1	D	38	THR	C-O	6.75	1.33	1.23
1	C	250	ARG	C-O	-5.94	1.17	1.24
1	C	242	HIS	C-O	-5.88	1.16	1.24
1	B	235	ARG	CA-C	5.60	1.60	1.52
1	D	54	ARG	C-N	5.59	1.41	1.33
1	B	246	ILE	CA-CB	5.56	1.57	1.53
1	C	252	THR	C-O	-5.54	1.17	1.24
1	D	97	GLN	C-O	-5.51	1.19	1.24
1	D	96	ALA	CA-CB	5.43	1.62	1.53
1	D	137	ILE	C-O	-5.39	1.18	1.24
1	A	99	HIS	C-O	-5.26	1.17	1.24
1	D	90	LEU	C-O	-5.14	1.17	1.23
1	D	54	ARG	N-CA	5.12	1.52	1.46

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	132	ILE	N-CA-C	5.13	115.25	110.42
1	B	170	LEU	N-CA-C	5.10	117.87	109.76
1	B	263	VAL	N-CA-C	5.02	115.74	110.62

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	43	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1900	0	1945	19	0
1	B	1831	0	1854	9	0
1	C	1800	0	1780	15	0
1	D	1792	0	1767	15	0
2	A	214	0	0	6	0
2	B	133	0	0	1	0
2	C	143	0	0	1	0
2	D	251	0	0	5	0
All	All	8064	0	7346	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:254:THR:HG22	2:A:413:HOH:O	1.48	1.10
1:B:130:SER:HB3	2:B:354:HOH:O	1.69	0.92
1:A:139:ILE:HD11	2:A:297:HOH:O	1.95	0.67
1:A:89:SER:HB3	1:D:138:THR:HG21	1.81	0.61
1:A:137:ILE:HD13	1:A:143:LEU:HD11	1.82	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:40:SER:O	1:B:43:ALA:HB2	2.01	0.59
1:D:251:VAL:HG21	2:D:297:HOH:O	2.03	0.58
1:D:126:ARG:NH2	2:D:302:HOH:O	2.38	0.57
1:B:42:VAL:HG21	1:B:57:LEU:HD11	1.86	0.56
1:A:117:ASP:CG	1:A:142:ARG:HH12	2.13	0.55
1:C:90:LEU:O	1:C:91:SER:HB3	2.06	0.55
1:C:260:LEU:C	1:C:262:THR:H	2.16	0.54
1:A:250:ARG:O	1:A:254:THR:HG23	2.08	0.53
1:D:57:LEU:O	1:D:61:VAL:HG23	2.08	0.53
1:B:20:LEU:HD13	1:C:23:GLY:HA3	1.92	0.52
1:C:52:THR:O	1:C:56:PHE:HD1	1.92	0.52
1:A:254:THR:HG21	2:A:335:HOH:O	2.10	0.51
1:A:135:VAL:HG12	1:C:133:MSE:HG3	1.93	0.51
1:A:81:GLU:CG	2:A:326:HOH:O	2.59	0.50
1:B:216:ILE:HD13	1:B:256:ILE:HG21	1.94	0.50
1:A:202:ASP:OD1	1:A:203:GLN:HG3	2.12	0.49
1:A:135:VAL:HG22	2:A:273:HOH:O	2.12	0.49
1:C:217:ARG:HD2	2:C:301:HOH:O	2.13	0.49
1:A:135:VAL:HG12	1:C:133:MSE:CG	2.43	0.48
1:C:199:CYS:O	1:C:212:MSE:HA	2.14	0.47
1:A:201:LEU:HD21	1:A:204:GLU:HB2	1.96	0.47
1:B:198:ILE:HD12	1:B:246:ILE:HG21	1.95	0.47
1:C:201:LEU:HD21	1:C:204:GLU:HB2	1.97	0.47
1:C:64:GLY:HA3	1:C:78:ARG:NH1	2.29	0.46
1:D:22:ARG:O	1:D:26:VAL:HG23	2.16	0.46
1:D:16:TYR:N	2:D:452:HOH:O	2.49	0.46
1:C:78:ARG:O	1:C:81:GLU:HG2	2.16	0.45
1:D:135:VAL:HG11	1:D:230:SER:CB	2.46	0.45
1:B:38:THR:O	1:B:42:VAL:HG23	2.16	0.45
1:A:81:GLU:HG2	2:A:326:HOH:O	2.16	0.45
1:C:198:ILE:HD11	1:C:249:LEU:HD23	1.99	0.45
1:A:76:THR:HB	1:A:77:PRO:HD2	2.00	0.43
1:D:161:ASP:HB3	2:D:448:HOH:O	2.17	0.43
1:D:228:ASN:OD1	1:D:228:ASN:C	2.61	0.43
1:A:35:GLN:HA	1:D:89:SER:HB2	2.01	0.42
1:B:87:LEU:HD23	1:B:90:LEU:HD11	2.01	0.42
1:C:64:GLY:HA3	1:C:78:ARG:HH11	1.83	0.42
1:A:228:ASN:OD1	1:A:228:ASN:C	2.61	0.42
1:D:199:CYS:O	1:D:212:MSE:HA	2.19	0.42
1:B:132:ILE:HG23	1:D:137:ILE:HG13	2.02	0.42
1:A:135:VAL:CG1	1:C:133:MSE:HG3	2.50	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:258:GLN:O	1:D:262:THR:HG23	2.20	0.41
1:C:198:ILE:HD12	1:C:214:ALA:HB2	2.02	0.41
1:D:181:THR:HG22	2:D:341:HOH:O	2.21	0.41
1:D:135:VAL:HG11	1:D:230:SER:HB2	2.03	0.41
1:A:145:ALA:O	1:A:151:GLY:HA3	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/265 (94%)	244 (98%)	4 (2%)	0	100	100
1	B	243/265 (92%)	236 (97%)	6 (2%)	1 (0%)	30	29
1	C	245/265 (92%)	234 (96%)	10 (4%)	1 (0%)	30	29
1	D	241/265 (91%)	236 (98%)	4 (2%)	1 (0%)	30	29
All	All	977/1060 (92%)	950 (97%)	24 (2%)	3 (0%)	37	37

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	261	ALA
1	D	53	ALA
1	B	42	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/217 (94%)	201 (98%)	3 (2%)	60	67
1	B	192/217 (88%)	186 (97%)	6 (3%)	35	39
1	C	185/217 (85%)	183 (99%)	2 (1%)	70	77
1	D	183/217 (84%)	179 (98%)	4 (2%)	47	53
All	All	764/868 (88%)	749 (98%)	15 (2%)	54	57

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	181	THR
1	A	212	MSE
1	A	222	LEU
1	B	52	THR
1	B	110[A]	SER
1	B	110[B]	SER
1	B	129	VAL
1	B	222	LEU
1	B	230	SER
1	C	161	ASP
1	C	201	LEU
1	D	126	ARG
1	D	191	LEU
1	D	222	LEU
1	D	228	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	242	HIS
1	A	264	ASN
1	B	105	HIS
1	D	105	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	247/265 (93%)	-0.04	3 (1%) 76 77	21, 33, 56, 69	0
1	B	243/265 (91%)	0.64	28 (11%) 11 11	19, 42, 95, 110	1 (0%)
1	C	244/265 (92%)	0.44	25 (10%) 13 14	24, 39, 87, 112	0
1	D	242/265 (91%)	0.30	36 (14%) 7 7	14, 27, 96, 124	0
All	All	976/1060 (92%)	0.33	92 (9%) 15 16	14, 36, 89, 124	1 (0%)

All (92) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	39	LEU	7.2
1	D	72	ALA	7.1
1	D	41	ASP	6.4
1	B	48	LEU	6.4
1	D	53	ALA	5.9
1	D	43	ALA	5.7
1	B	17	VAL	5.6
1	D	38	THR	5.2
1	D	73	PHE	5.0
1	D	49	THR	4.8
1	D	67	ALA	4.8
1	D	57	LEU	4.5
1	C	49	THR	4.3
1	D	44	ARG	4.3
1	B	44	ARG	4.2
1	D	40	SER	4.2
1	B	181	THR	4.2
1	B	38	THR	4.1
1	D	42	VAL	4.0
1	D	51	ALA	4.0
1	D	54	ARG	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	39	LEU	3.9
1	A	219	ALA	3.9
1	D	61	VAL	3.9
1	D	46	THR	3.9
1	C	48	LEU	3.9
1	D	50	ARG	3.8
1	D	45	ALA	3.8
1	B	43	ALA	3.8
1	C	53	ALA	3.7
1	C	17	VAL	3.7
1	C	21	ALA	3.6
1	D	48	LEU	3.6
1	B	32	HIS	3.5
1	D	74	TRP	3.5
1	C	41	ASP	3.4
1	C	70	GLY	3.4
1	C	52	THR	3.3
1	C	40	SER	3.3
1	C	51	ALA	3.2
1	D	32	HIS	3.2
1	B	220	SER	3.1
1	B	51	ALA	3.1
1	C	47	ASP	3.1
1	D	52	THR	3.1
1	C	43	ALA	3.0
1	B	49	THR	3.0
1	B	56	PHE	2.9
1	B	39	LEU	2.9
1	D	58	LEU	2.9
1	C	42	VAL	2.9
1	B	172	ILE	2.8
1	C	71	SER	2.8
1	B	70	GLY	2.7
1	A	181	THR	2.7
1	B	219	ALA	2.7
1	D	35	GLN	2.7
1	B	58	LEU	2.7
1	C	44	ARG	2.7
1	D	37	ARG	2.7
1	B	36	ARG	2.6
1	B	180	ILE	2.6
1	D	33	ARG	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	40	SER	2.6
1	B	20	LEU	2.6
1	C	50	ARG	2.5
1	C	57	LEU	2.5
1	B	33	ARG	2.5
1	C	58	LEU	2.5
1	D	66	VAL	2.4
1	C	73	PHE	2.4
1	C	27	ILE	2.4
1	B	69	ASP	2.4
1	B	53	ALA	2.4
1	B	72	ALA	2.4
1	D	36	ARG	2.4
1	D	62	GLU	2.3
1	C	38	THR	2.3
1	B	71	SER	2.3
1	A	135	VAL	2.2
1	B	50	ARG	2.2
1	D	55	ARG	2.2
1	B	52	THR	2.2
1	C	261	ALA	2.2
1	D	56	PHE	2.2
1	C	60	LEU	2.2
1	D	47	ASP	2.2
1	D	30	PHE	2.1
1	C	72	ALA	2.1
1	B	42	VAL	2.1
1	D	29	CYS	2.1
1	D	60	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.