



Full wwPDB X-ray Structure Validation Report i

Nov 12, 2024 – 11:48 AM EST

PDB ID : 3IXJ
Title : Crystal structure of beta-secretase 1 in complex with selective beta-secretase 1 inhibitor
Authors : Borkakoti, N.; Lindberg, J.; Nystrom, S.
Deposited on : 2009-09-04
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)


There are 4 unique types of molecules in this entry. The entry contains 9241 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	373	2940	1882	489	555	14	0	0	0
1	B	373	2940	1882	489	555	14	0	0	0
1	C	371	2925	1874	486	551	14	0	0	0

- Molecule 2 is N-[4-(1-BENZYLCARBAMOYL-2-METHYL-PROPYLCARBAMOYL)-1-(3,5-DIFLUOROPHENOXYMETHYL)-2-HYDROXY-4-METHOXY-BUTYL]-5-(METHANESULFONYL-METHYL-AMINO)-N'-(1-PHENYLETHYL)-ISOPHTHALAMIDE (three-letter code: 586) (formula: C₄₃H₅₁F₂N₅O₉S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O	S	
2	A	1	60	43	2	5	9	1	0
2	B	1	60	43	2	5	9	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	F	N	O	S		
2	C	1	60	43	2	5	9	1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	A	1	5	4	1	0	0
3	B	1	5	4	1	0	0
3	C	1	5	4	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	82	82	82	0	0
4	B	89	89	89	0	0
4	C	70	70	70	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-secretase 1



- Molecule 1: Beta-secretase 1



- Molecule 1: Beta-secretase 1



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.91Å 102.70Å 100.42Å 90.00° 103.42° 90.00°	Depositor
Resolution (Å)	62.99 – 2.20 62.99 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.5 (62.99-2.20) 99.5 (62.99-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$< I/\sigma(I) >$ ¹	3.98 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R , R_{free}	0.212 , 0.243 0.211 , 0.242	Depositor DCC
R_{free} test set	4101 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	19.3	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 30.9	EDS
L-test for twinning ²	$< L > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9241	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 586, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/3014	0.57	0/4092
1	B	0.55	0/3014	0.57	0/4092
1	C	0.53	0/2999	0.56	0/4072
All	All	0.54	0/9027	0.57	0/12256

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2940	0	2849	20	0
1	B	2940	0	2849	9	0
1	C	2925	0	2838	19	0
2	A	60	0	51	0	0
2	B	60	0	51	0	0
2	C	60	0	51	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
4	A	82	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	89	0	0	0	0
4	C	70	0	0	0	0
All	All	9241	0	8689	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:305:PHE:HD1	1:A:316:VAL:HG11	1.41	0.84
1:C:424:THR:HB	1:C:427:MET:HE2	1.61	0.82
1:B:305:PHE:HD1	1:B:316:VAL:HG11	1.56	0.70
1:C:424:THR:HB	1:C:427:MET:CE	2.23	0.69
1:C:252:VAL:HG11	1:C:427:MET:HG2	1.77	0.67
1:A:48:VAL:CG1	1:B:125:GLU:HB2	2.25	0.67
1:A:305:PHE:CD1	1:A:316:VAL:HG11	2.28	0.65
1:B:319:GLN:CD	1:B:319:GLN:H	2.00	0.64
1:A:116:TYR:HE2	1:A:123:LYS:HB2	1.63	0.64
1:C:349:LEU:H	1:C:352:GLN:NE2	1.97	0.62
1:C:408:HIS:CE1	1:C:416:ALA:H	2.17	0.62
1:B:242:ARG:HD2	1:B:250:ILE:HD11	1.81	0.60
1:A:308:GLY:HA3	1:A:314:GLN:HE21	1.66	0.60
1:C:349:LEU:H	1:C:352:GLN:HE21	1.50	0.60
1:C:342:GLN:C	1:C:427:MET:HE1	2.22	0.60
1:A:116:TYR:CE2	1:A:123:LYS:HB2	2.39	0.57
1:C:317:CYS:HG	1:C:367:CYS:HG	1.52	0.56
1:A:116:TYR:OH	1:A:118:PRO:HB3	2.07	0.54
1:A:306:PRO:O	1:A:309:PHE:HB3	2.07	0.54
1:C:365:ASP:OD1	1:C:365:ASP:N	2.42	0.52
1:A:116:TYR:CE2	1:A:123:LYS:HD3	2.44	0.52
1:A:48:VAL:HG12	1:B:125:GLU:HB2	1.92	0.51
1:B:309:PHE:CE1	1:B:316:VAL:HG22	2.47	0.50
1:A:57:LYS:N	1:A:57:LYS:HE2	2.29	0.47
1:C:270:TYR:HA	1:C:271:ASP:HA	1.70	0.47
1:C:66:MET:HG2	1:C:135:ILE:HG12	1.97	0.47
1:B:346:ILE:HG22	1:B:418:VAL:HG22	1.97	0.46
1:B:270:TYR:HA	1:B:271:ASP:HA	1.70	0.45
1:C:343:SER:HB3	1:C:427:MET:CE	2.47	0.44
1:A:307:ASP:HA	1:A:310:TRP:HD1	1.82	0.43
1:A:62:TYR:CE2	1:A:218:VAL:HG13	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:117:VAL:HG22	1:C:176:ARG:HB2	2.01	0.43
1:A:116:TYR:HE2	1:A:123:LYS:HD3	1.83	0.43
1:C:405:SER:O	1:C:408:HIS:HD2	2.02	0.42
1:C:408:HIS:HE1	1:C:416:ALA:H	1.65	0.42
1:A:293:VAL:O	1:A:297:LYS:HB2	2.19	0.42
1:C:68:VAL:HG12	1:C:133:VAL:HG22	2.00	0.42
1:A:309:PHE:CE1	1:A:316:VAL:HG22	2.54	0.42
1:B:227:ILE:HG23	1:B:390:TYR:HE2	1.84	0.41
1:A:392:VAL:O	1:A:400:ILE:HA	2.20	0.41
1:C:324:PRO:O	1:C:327:ILE:HG12	2.19	0.41
1:A:270:TYR:HA	1:A:271:ASP:HA	1.72	0.41
1:C:319:GLN:HE21	1:C:319:GLN:HB2	1.66	0.41
1:A:253:ARG:HB3	1:A:334:TYR:HB2	2.03	0.41
1:C:63:TYR:CD1	1:C:76:ASN:HB3	2.56	0.41
1:A:276:ASP:O	1:A:382:GLY:HA2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	367/388 (95%)	360 (98%)	7 (2%)	0	100 100
1	B	367/388 (95%)	360 (98%)	7 (2%)	0	100 100
1	C	365/388 (94%)	358 (98%)	6 (2%)	1 (0%)	37 42
All	All	1099/1164 (94%)	1078 (98%)	20 (2%)	1 (0%)	48 57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	320	ALA

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	319/331 (96%)	312 (98%)	7 (2%)	47 61
1	B	319/331 (96%)	313 (98%)	6 (2%)	52 67
1	C	317/331 (96%)	305 (96%)	12 (4%)	28 37
All	All	955/993 (96%)	930 (97%)	25 (3%)	41 54

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	LYS
1	A	218	VAL
1	A	245	TRP
1	A	302	THR
1	A	408	HIS
1	A	409	VAL
1	A	423	VAL
1	B	245	TRP
1	B	287	LYS
1	B	319	GLN
1	B	358	GLU
1	B	404	VAL
1	B	423	VAL
1	C	48	VAL
1	C	98	ARG
1	C	112	ARG
1	C	121	GLN
1	C	245	TRP
1	C	259	GLN
1	C	304	LYS
1	C	307	ASP
1	C	319	GLN
1	C	365	ASP
1	C	366	ASP
1	C	412	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	ASN
1	A	314	GLN
1	A	341	ASN
1	A	352	GLN
1	B	76	ASN
1	B	314	GLN
1	B	319	GLN
1	B	364	GLN
1	B	374	GLN
1	C	76	ASN
1	C	121	GLN
1	C	259	GLN
1	C	319	GLN
1	C	341	ASN
1	C	352	GLN
1	C	374	GLN
1	C	408	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	700	-	4,4,4	0.27	0	6,6,6	0.11	0
2	586	B	500	-	62,63,63	1.11	2 (3%)	83,88,88	1.38	11 (13%)
2	586	A	500	-	62,63,63	1.01	2 (3%)	83,88,88	1.35	13 (15%)
3	SO4	C	700	-	4,4,4	0.22	0	6,6,6	0.16	0
2	586	C	500	-	62,63,63	1.03	2 (3%)	83,88,88	1.29	8 (9%)
3	SO4	B	700	-	4,4,4	0.22	0	6,6,6	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	586	B	500	-	-	12/66/66/66	0/4/4/4
2	586	C	500	-	-	6/66/66/66	0/4/4/4
2	586	A	500	-	-	11/66/66/66	0/4/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	586	S-N5	-5.18	1.55	1.64
2	C	500	586	S-N5	-4.98	1.55	1.64
2	A	500	586	S-N5	-4.63	1.56	1.64
2	B	500	586	C44-S	-3.85	1.67	1.75
2	A	500	586	C44-S	-3.47	1.68	1.75
2	C	500	586	C44-S	-3.33	1.68	1.75

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	586	O7-S-O8	-5.59	110.10	118.46
2	C	500	586	O7-S-O8	-5.16	110.74	118.46
2	C	500	586	O7-S-N5	5.05	112.92	107.14
2	B	500	586	O7-S-O8	-5.00	110.98	118.46
2	A	500	586	O2-C6-C5	3.90	114.63	108.13
2	C	500	586	O2-C6-C5	3.84	114.53	108.13
2	B	500	586	O2-C6-C5	3.63	114.19	108.13
2	B	500	586	C21-C22-C23	3.36	121.75	117.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	586	C10-C8-C9	-3.36	103.19	111.38
2	B	500	586	C22-C23-C24	-3.14	119.69	123.50
2	B	500	586	O8-S-N5	3.13	110.72	107.14
2	A	500	586	C22-C23-C24	-3.08	119.76	123.50
2	A	500	586	O7-S-N5	3.01	110.59	107.14
2	B	500	586	O7-S-N5	2.96	110.53	107.14
2	C	500	586	C22-C23-C24	-2.68	120.25	123.50
2	A	500	586	C21-C22-C23	2.54	120.72	117.55
2	C	500	586	C44-S-N5	2.49	109.86	106.82
2	A	500	586	C23-C24-C25	2.44	119.64	116.08
2	C	500	586	C1-C5-N1	2.41	114.49	109.97
2	A	500	586	C6-O2-C21	2.32	122.96	117.85
2	B	500	586	C10-C8-N2	2.31	117.03	111.44
2	B	500	586	C1-C5-N1	2.30	114.28	109.97
2	A	500	586	C1-C5-N1	2.21	114.12	109.97
2	A	500	586	C44-S-N5	2.17	109.48	106.82
2	C	500	586	C23-C24-C25	2.17	119.24	116.08
2	A	500	586	O8-S-N5	2.15	109.60	107.14
2	B	500	586	C11-C10-C8	2.12	116.98	111.16
2	C	500	586	C21-C22-C23	2.11	120.19	117.55
2	A	500	586	C14-C13-N3	-2.10	108.65	113.07
2	A	500	586	C10-C8-C9	-2.09	106.27	111.38
2	A	500	586	C24-C25-C26	-2.06	121.00	123.50
2	B	500	586	C21-C26-C25	2.05	120.11	117.55

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500	586	C31-C32-N5-S
2	A	500	586	C33-C32-N5-S
2	A	500	586	N1-C5-C6-O2
2	B	500	586	C31-C32-N5-S
2	B	500	586	C33-C32-N5-S
2	B	500	586	N1-C5-C6-O2
2	C	500	586	C31-C32-N5-S
2	C	500	586	C33-C32-N5-S
2	C	500	586	N1-C5-C6-O2
2	A	500	586	C12-C10-C8-C9
2	B	500	586	C12-C10-C8-C9
2	B	500	586	C12-C10-C8-N2
2	A	500	586	C12-C10-C8-N2

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	500	586	C11-C10-C8-N2
2	B	500	586	C11-C10-C8-C9
2	A	500	586	C11-C10-C8-C9
2	A	500	586	C11-C10-C8-N2
2	A	500	586	C5-C6-O2-C21
2	C	500	586	C5-C6-O2-C21
2	A	500	586	C1-C5-C6-O2
2	B	500	586	C1-C5-C6-O2
2	C	500	586	C1-C5-C6-O2
2	B	500	586	C5-C6-O2-C21
2	B	500	586	C33-C27-C28-O5
2	C	500	586	O9-C2-C4-N2
2	B	500	586	C29-C27-C28-O5
2	B	500	586	C33-C27-C28-N1
2	A	500	586	C33-C27-C28-O5
2	A	500	586	C29-C27-C28-O5


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	373/388 (96%)	0.13	26 (6%) 24 21	5, 17, 38, 56	0
1	B	373/388 (96%)	0.09	22 (5%) 29 26	5, 18, 38, 53	0
1	C	371/388 (95%)	0.19	26 (7%) 24 21	4, 19, 47, 57	0
All	All	1117/1164 (95%)	0.14	74 (6%) 26 23	4, 18, 42, 57	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	205	ALA	5.8
1	A	306	PRO	5.4
1	B	410	HIS	5.3
1	A	412	GLU	5.1
1	A	409	VAL	5.0
1	A	116	TYR	5.0
1	C	205	ALA	5.0
1	B	306	PRO	4.9
1	A	363	SER	4.5
1	B	304	LYS	4.2
1	B	217	SER	4.2
1	C	306	PRO	4.2
1	B	363	SER	4.2
1	C	46	SER	4.2
1	C	365	ASP	4.1
1	A	358	GLU	4.1
1	A	364	GLN	4.1
1	C	366	ASP	4.0
1	B	412	GLU	3.9
1	A	413	PHE	3.8
1	A	410	HIS	3.8
1	A	217	SER	3.8
1	C	358	GLU	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	46	SER	3.7
1	A	58	SER	3.6
1	B	302	THR	3.5
1	B	364	GLN	3.5
1	B	413	PHE	3.5
1	B	319	GLN	3.4
1	B	205	ALA	3.4
1	C	301	SER	3.4
1	B	320	ALA	3.3
1	C	299	ALA	3.3
1	A	307	ASP	3.3
1	C	319	GLN	3.3
1	C	314	GLN	3.1
1	A	302	THR	3.1
1	C	304	LYS	3.1
1	C	315	LEU	3.1
1	C	310	TRP	3.0
1	C	305	PHE	2.9
1	C	112	ARG	2.8
1	A	57	LYS	2.8
1	A	304	LYS	2.8
1	C	300	SER	2.8
1	C	318	TRP	2.8
1	B	52	ASP	2.7
1	A	59	GLY	2.7
1	B	307	ASP	2.7
1	A	408	HIS	2.7
1	C	307	ASP	2.7
1	C	302	THR	2.7
1	B	358	GLU	2.7
1	C	298	ALA	2.6
1	A	218	VAL	2.6
1	B	409	VAL	2.6
1	C	311	LEU	2.6
1	A	314	GLN	2.6
1	C	320	ALA	2.6
1	B	315	LEU	2.6
1	A	433	ASN	2.6
1	C	367	CYS	2.6
1	C	326	ASN	2.5
1	A	319	GLN	2.5
1	C	317	CYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	411	ASP	2.4
1	B	97	HIS	2.4
1	A	112	ARG	2.4
1	A	428	GLU	2.3
1	B	316	VAL	2.3
1	C	297	LYS	2.2
1	B	321	GLY	2.1
1	A	320	ALA	2.1
1	A	411	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	700	5/5	0.85	0.15	64,64,65,65	0
3	SO4	B	700	5/5	0.88	0.20	57,57,58,58	0
3	SO4	C	700	5/5	0.91	0.15	52,53,53,54	0
2	586	A	500	60/60	0.94	0.08	7,12,23,25	0
2	586	C	500	60/60	0.96	0.07	5,11,23,24	0
2	586	B	500	60/60	0.97	0.06	4,9,19,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.