



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 3, 2025 – 07:08 pm BST

PDB ID : 5L70 / pdb_00005L70
Title : CRYSTAL STRUCTURE OF HUMAN CARBONIC ANHYDRASE II IN
COMPLEX WITH A QUINOLINE OLIGOAMIDE FOLDAMER
Authors : Vallade, M.; Langlois d'Estaintot, B.; Granier, T.; Huc, I.
Deposited on : 2016-06-01
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	<div> <div>%</div> <div> <div></div> <div>93%</div> <div>6%</div> <div>.</div> </div> </div>
1	B	260	<div> <div>3%</div> <div> <div></div> <div>96%</div> <div>.</div> <div>.</div> </div> </div>
2	C	6	<div> <div>17%</div> <div>67%</div> <div>17%</div> </div>
2	D	6	<div> <div>17%</div> <div>83%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4593 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbonic anhydrase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	258	Total	C	N	O	S	0	3	0
			2035	1308	343	382	2			
1	B	258	Total	C	N	O	S	0	0	0
			1989	1280	340	367	2			

- Molecule 2 is a protein called Aromatic foldamer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	6	Total	C	N	O	S	0	0	0
			99	68	12	18	1			
2	D	6	Total	C	N	O	S	0	0	0
			99	68	12	18	1			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	9	Total	Zn	0	0
			9	9		
3	B	4	Total	Zn	0	0
			4	4		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	189	Total	O	0	0
			189	189		
5	B	131	Total	O	0	0
			131	131		
5	C	5	Total	O	0	0
			5	5		
5	D	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Carbonic anhydrase 2




- Molecule 1: Carbonic anhydrase 2



- Molecule 2: Aromatic foldamer



- Molecule 2: Aromatic foldamer



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.21Å 83.37Å 79.41Å 90.00° 99.98° 90.00°	Depositor
Resolution (Å)	78.21 – 2.20 78.21 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.0 (78.21-2.20) 96.0 (78.21-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.94 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.174 , 0.238 0.181 , 0.236	Depositor DCC
R_{free} test set	1402 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	23.7	Xtriage
Anisotropy	0.012	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4593	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: QOL, QVE, QUK, A1IJ4, ZN, QVS, 4SO, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/2106	0.90	0/2869
1	B	0.78	0/2051	0.89	0/2800
All	All	0.82	0/4157	0.90	0/5669

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	303	QOL	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2035	0	1912	9	0
1	B	1989	0	1852	4	0
2	C	99	0	6	0	0
2	D	99	0	6	0	0
3	A	9	0	0	0	0
3	B	4	0	0	0	0
4	A	12	0	16	0	0
4	B	12	0	16	0	0
5	A	189	0	0	2	0
5	B	131	0	0	1	0
5	C	5	0	0	0	0
5	D	9	0	0	1	0
All	All	4593	0	3808	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252[B]:ASN:O	1:A:252[B]:ASN:CG	2.32	0.72
1:A:10:HIS:CD2	5:A:555:HOH:O	2.53	0.60
1:B:157:GLN:NE2	5:B:401:HOH:O	2.34	0.59
1:A:189:ASP:HB2	1:A:212:LYS:HE3	1.86	0.58
1:B:133:ALA:O	1:B:139:GLY:HA3	2.06	0.56
1:A:133:ALA:O	1:A:139:GLY:HA3	2.06	0.55
1:A:167:LYS:NZ	5:A:402:HOH:O	2.42	0.53
1:A:3:HIS:CE1	5:D:404:HOH:O	2.70	0.44
1:B:89:ARG:O	1:B:122:HIS:HA	2.17	0.44
1:A:45:LYS:O	1:A:82:GLY:HA2	2.18	0.43
1:A:89:ARG:O	1:A:122:HIS:HA	2.18	0.43
1:A:116:ALA:HB3	1:A:147:LEU:HB2	2.01	0.42
1:B:45:LYS:O	1:B:82:GLY:HA2	2.19	0.42

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/260 (100%)	248 (96%)	11 (4%)	0	100	100
1	B	256/260 (98%)	248 (97%)	8 (3%)	0	100	100
All	All	515/520 (99%)	496 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/225 (94%)	210 (100%)	1 (0%)	86	93
1	B	201/225 (89%)	200 (100%)	1 (0%)	86	93
All	All	412/450 (92%)	410 (100%)	2 (0%)	86	93

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	216	SER
1	B	216	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	36	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	QUK	C	305	2	19,19,20	1.87	2 (10%)	22,25,27	2.71	7 (31%)
2	QVS	D	304	2	15,15,16	1.72	2 (13%)	19,21,23	3.00	5 (26%)
2	QVE	C	306	2,3	20,20,20	1.74	1 (5%)	27,28,28	1.66	8 (29%)
2	QVS	C	304	2	15,15,16	2.05	2 (13%)	19,21,23	2.87	6 (31%)
2	A1IJ4	C	302	2	16,16,17	0.88	1 (6%)	17,18,20	1.48	1 (5%)
2	QOL	D	303	2	21,21,22	2.18	1 (4%)	23,28,30	2.44	6 (26%)
2	QUK	D	305	2	19,19,20	1.77	1 (5%)	22,25,27	2.81	7 (31%)
2	QVE	D	306	2,3	20,20,20	1.79	1 (5%)	27,28,28	2.03	8 (29%)
2	A1IJ4	D	302	2	16,16,17	0.95	1 (6%)	17,18,20	1.28	1 (5%)
2	QOL	C	303	2	21,21,22	1.88	1 (4%)	23,28,30	2.82	5 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QUK	C	305	2	-	2/7/7/9	0/2/2/2
2	QVS	D	304	2	-	2/2/2/4	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QVE	C	306	2,3	-	2/9/9/9	0/2/2/2
2	QVS	C	304	2	-	2/2/2/4	0/2/2/2
2	A1IJ4	C	302	2	-	1/10/10/11	0/1/1/1
2	QOL	D	303	2	-	4/11/11/13	0/2/2/2
2	QUK	D	305	2	-	2/7/7/9	0/2/2/2
2	QVE	D	306	2,3	-	6/9/9/9	0/2/2/2
2	A1IJ4	D	302	2	-	1/10/10/11	0/1/1/1
2	QOL	C	303	2	-	2/11/11/13	0/2/2/2

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	303	QOL	C10-C	-9.30	1.38	1.48
2	C	303	QOL	C10-C	-7.49	1.40	1.48
2	C	305	QUK	C10-C	-7.22	1.41	1.48
2	D	305	QUK	C10-C	-6.89	1.41	1.48
2	D	306	QVE	CA-C	-6.83	1.41	1.50
2	C	306	QVE	CA-C	-6.63	1.41	1.50
2	C	304	QVS	CA-C	-6.55	1.41	1.48
2	D	304	QVS	CA-C	-5.16	1.43	1.48
2	C	304	QVS	CA-N11	3.03	1.36	1.33
2	D	302	A1IJ4	C128-C127	-2.95	1.41	1.51
2	D	304	QVS	CA-N11	2.85	1.36	1.33
2	C	302	A1IJ4	C128-C127	-2.78	1.41	1.51
2	C	305	QUK	C10-N11	2.12	1.35	1.33

All (54) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	304	QVS	CA-N11-C7	9.77	125.52	118.11
2	D	305	QUK	C10-N11-C7	9.47	125.29	118.11
2	C	304	QVS	CA-N11-C7	9.46	125.28	118.11
2	C	305	QUK	C10-N11-C7	8.90	124.86	118.11
2	C	303	QOL	C-C10-N11	8.29	122.78	114.66
2	C	303	QOL	C10-N11-C7	7.90	124.10	118.11
2	D	303	QOL	C-C10-N11	6.64	121.16	114.66
2	D	303	QOL	C10-N11-C7	6.16	122.78	118.11
2	C	305	QUK	O-C-C10	-5.44	119.07	124.22
2	D	306	QVE	CG-OB-C8	5.43	123.13	116.95
2	C	302	A1IJ4	O-C-N31	-5.32	119.52	124.89
2	D	305	QUK	C9-C10-C	4.88	125.44	121.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	304	QVS	O-C-CA	-4.83	119.64	124.22
2	C	304	QVS	O-C-CA	-4.83	119.65	124.22
2	D	303	QOL	C3-CA-N	4.64	129.64	120.36
2	D	302	A1IJ4	O-C-N31	-4.61	120.24	124.89
2	D	305	QUK	O-C-C10	-4.54	119.92	124.22
2	C	303	QOL	C3-CA-N	4.50	129.35	120.36
2	D	306	QVE	C-CA-N11	4.49	123.45	116.28
2	C	305	QUK	C9-C10-C	4.17	124.83	121.23
2	D	304	QVS	C-CA-N11	4.16	118.74	114.66
2	C	303	QOL	C7-CA-N	-3.62	111.17	118.07
2	C	306	QVE	CA-N11-C7	3.60	124.82	117.24
2	D	304	QVS	C9-CA-C	3.52	124.27	121.23
2	D	303	QOL	C7-CA-N	-3.51	111.37	118.07
2	C	304	QVS	C-CA-N11	3.45	118.05	114.66
2	C	306	QVE	CG-OB-C8	3.45	120.88	116.95
2	D	305	QUK	C3-CA-N	3.41	127.18	120.36
2	D	306	QVE	C2-C7-N11	3.40	122.00	118.64
2	C	304	QVS	C9-CA-C	3.18	123.98	121.23
2	C	305	QUK	C3-CA-N	3.15	126.65	120.36
2	D	304	QVS	C3-C2-N	3.10	126.56	120.36
2	D	306	QVE	CA-N11-C7	3.08	123.72	117.24
2	C	304	QVS	C3-C2-N	3.04	126.45	120.36
2	C	306	QVE	C2-C7-N11	2.98	121.59	118.64
2	D	306	QVE	C9-CA-N11	-2.81	117.65	124.50
2	C	306	QVE	C-CA-N11	2.79	120.73	116.28
2	C	306	QVE	C9-CA-N11	-2.74	117.82	124.50
2	D	305	QUK	C3-CA-C7	-2.65	115.89	120.06
2	D	306	QVE	OXT-C-O	-2.59	117.60	123.35
2	D	305	QUK	CG-OB-C8	2.57	125.42	117.74
2	C	303	QOL	OB-CG-CD	2.49	114.71	108.21
2	D	303	QOL	OB-CG-CD	2.47	114.66	108.21
2	D	303	QOL	O-C-C10	-2.45	121.90	124.22
2	D	305	QUK	C6-C7-N11	-2.37	117.89	122.78
2	D	306	QVE	OXT-C-CA	2.33	120.00	114.69
2	C	305	QUK	CG-OB-C8	2.28	124.56	117.74
2	C	306	QVE	OXT-C-CA	2.27	119.85	114.69
2	C	306	QVE	OXT-C-O	-2.26	118.33	123.35
2	C	305	QUK	C6-C7-N11	-2.22	118.21	122.78
2	D	306	QVE	C3-C2-N	2.19	124.74	120.36
2	C	306	QVE	C6-C7-N11	-2.15	118.34	122.78
2	C	304	QVS	C6-C7-N11	-2.09	118.47	122.78
2	C	305	QUK	C3-CA-C7	-2.03	116.87	120.06

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	303	QOL	CE2-CD-CG-OB
2	D	303	QOL	CE1-CD-CG-OB
2	D	303	QOL	CE2-CD-CG-OB
2	D	303	QOL	O-C-C10-C9
2	D	303	QOL	O-C-C10-N11
2	C	304	QVS	O-C-CA-C9
2	C	304	QVS	O-C-CA-N11
2	D	304	QVS	O-C-CA-C9
2	D	304	QVS	O-C-CA-N11
2	D	306	QVE	OE2-CD-CG-OB
2	D	302	A1IJ4	C119-C120-C121-O71
2	D	306	QVE	OE1-CD-CG-OB
2	C	305	QUK	CG-CD-CE-N1
2	C	302	A1IJ4	C119-C120-C121-O71
2	C	303	QOL	CE1-CD-CG-OB
2	C	306	QVE	OE1-CD-CG-OB
2	D	306	QVE	C9-C8-OB-CG
2	C	306	QVE	OE2-CD-CG-OB
2	D	306	QVE	O-C-CA-C9
2	D	305	QUK	CG-CD-CE-N1
2	D	306	QVE	O-C-CA-N11
2	D	306	QVE	C6-C8-OB-CG
2	C	305	QUK	CE-CD-CG-OB
2	D	305	QUK	CD-CG-OB-C8

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 13 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	B	303	-	5,5,5	0.31	0	5,5,5	0.47	0
4	GOL	B	306	-	5,5,5	0.36	0	5,5,5	0.26	0
4	GOL	A	311	-	5,5,5	0.35	0	5,5,5	0.40	0
4	GOL	A	303	-	5,5,5	0.34	0	5,5,5	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	303	-	-	4/4/4/4	-
4	GOL	B	306	-	-	2/4/4/4	-
4	GOL	A	311	-	-	2/4/4/4	-
4	GOL	A	303	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	311	GOL	O1-C1-C2-C3
4	B	303	GOL	O1-C1-C2-C3
4	B	303	GOL	C1-C2-C3-O3
4	B	306	GOL	O1-C1-C2-C3
4	A	303	GOL	O1-C1-C2-C3
4	A	311	GOL	O1-C1-C2-O2
4	B	303	GOL	O2-C2-C3-O3
4	B	306	GOL	O1-C1-C2-O2
4	A	303	GOL	O1-C1-C2-O2
4	B	303	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	258/260 (99%)	-0.38	3 (1%) 76 73	12, 26, 43, 60	3 (1%)
1	B	258/260 (99%)	0.21	7 (2%) 56 53	16, 37, 64, 90	0
2	C	0/6	-	-	-	-
2	D	0/6	-	-	-	-
All	All	516/532 (96%)	-0.08	10 (1%) 66 62	12, 31, 59, 90	3 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	HIS	2.7
1	A	252[A]	ASN	2.6
1	B	152	ALA	2.5
1	B	161	ASP	2.4
1	A	10	HIS	2.4
1	B	3	HIS	2.3
1	B	236	PRO	2.3
1	B	228	LEU	2.2
1	B	168	THR	2.1
1	B	234	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	QVS	C	304	14/15	0.91	0.08	22,25,28,32	0
2	QOL	C	303	20/21	0.92	0.09	17,24,49,55	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	QUK	D	305	18/19	0.92	0.09	19,24,44,44	0
2	QVE	C	306	19/19	0.92	0.10	20,26,43,60	0
2	QVE	D	306	19/19	0.92	0.09	18,23,41,48	0
2	QUK	C	305	18/19	0.93	0.09	23,30,49,51	0
2	A1IJ4	D	302	16/17	0.93	0.08	15,26,33,34	0
2	QOL	D	303	20/21	0.94	0.08	16,22,40,43	0
2	A1IJ4	C	302	16/17	0.96	0.07	14,22,40,41	0
2	QVS	D	304	14/15	0.96	0.06	12,20,28,28	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	B	306	6/6	0.89	0.10	37,47,52,52	0
4	GOL	B	303	6/6	0.91	0.16	21,33,36,44	0
3	ZN	A	307	1/1	0.92	0.08	69,69,69,69	0
3	ZN	B	304	1/1	0.92	0.08	63,63,63,63	0
3	ZN	A	310	1/1	0.93	0.08	34,34,34,34	1
4	GOL	A	311	6/6	0.93	0.07	25,26,35,38	0
4	GOL	A	303	6/6	0.94	0.13	19,30,37,51	0
3	ZN	A	308	1/1	0.95	0.09	58,58,58,58	0
3	ZN	A	304	1/1	0.96	0.06	56,56,56,56	0
3	ZN	B	305	1/1	0.96	0.07	31,31,31,31	1
3	ZN	A	309	1/1	0.98	0.05	47,47,47,47	0
3	ZN	A	302	1/1	0.99	0.03	20,20,20,20	0
3	ZN	B	302	1/1	0.99	0.02	27,27,27,27	0
3	ZN	A	305	1/1	0.99	0.03	30,30,30,30	0
3	ZN	A	306	1/1	0.99	0.03	26,26,26,26	0
3	ZN	B	301	1/1	1.00	0.01	18,18,18,18	0
3	ZN	A	301	1/1	1.00	0.01	12,12,12,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.