



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 1, 2025 – 10:03 AM EDT

PDB ID : 9NGT / pdb_00009ngt
Title : Crystal structure of CRBN-DDB1 and FPFT-2216 in complex with mTOR
Authors : Petzold, G.; Trenh, P.; Bunker, R.D.; Tsai, J.H.C.
Deposited on : 2025-02-22
Resolution : 2.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1044 (2.98-2.94)
Clashscore	180529	1097 (2.98-2.94)
Ramachandran outliers	177936	1049 (2.98-2.94)
Sidechain outliers	177891	1049 (2.98-2.94)
RSRZ outliers	164620	1044 (2.98-2.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1140	<div> <div>0% (red)</div> <div>67% (green)</div> <div>29% (grey)</div> </div>
2	B	442	<div> <div>4% (red)</div> <div>75% (green)</div> <div>5% (yellow)</div> <div>20% (grey)</div> </div>
3	C	98	<div> <div>2% (red)</div> <div>87% (green)</div> <div>7% (yellow)</div> <div>5% (grey)</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10019 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA damage-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	807	Total	C	N	O	S	0	0	0
			6345	4023	1068	1220	34			

- Molecule 2 is a protein called Protein cereblon.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	355	Total	C	N	O	S	0	0	0
			2859	1828	488	520	23			

- Molecule 3 is a protein called Serine/threonine-protein kinase mTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	93	Total	C	N	O	S	0	0	0
			794	505	140	142	7			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	2017	GLY	-	expression tag	UNP P42345


- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		


- Molecule 5 is (3S)-3-[(4M)-4-(4-methoxythiophen-3-yl)-1H-1,2,3-triazol-1-yl]piperidine-2,6-dione (CCD ID: A1BC8) (formula: C₁₂H₁₂N₄O₃S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
5	B	1	20	12	4	3	1	0	0



● Molecule 3: Serine/threonine-protein kinase mTOR



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	183.18Å 81.21Å 99.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.78 – 2.95 43.78 – 2.95	Depositor EDS
% Data completeness (in resolution range)	93.1 (43.78-2.95) 93.0 (43.78-2.95)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 2.61Å)	Xtriage
Refinement program	PHENIX dev_5430	Depositor
R, R_{free}	0.231 , 0.260 0.233 , 0.263	Depositor DCC
R_{free} test set	3738 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å ²)	68.0	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	10019	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1BC8, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/6459	0.87	1/8736 (0.0%)
2	B	0.70	1/2924 (0.0%)	0.85	1/3964 (0.0%)
3	C	0.64	0/816	1.02	0/1094
All	All	0.65	1/10199 (0.0%)	0.87	2/13794 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	162	ARG	CA-C	5.44	1.56	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1106	GLN	CB-CG-CD	6.31	123.32	112.60
2	B	186	PRO	N-CA-C	5.03	119.20	111.21

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6345	0	6319	11	0
2	B	2859	0	2867	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	794	0	756	2	0
4	B	1	0	0	0	0
5	B	20	0	0	2	0
All	All	10019	0	9942	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:MET:HE2	1:A:169:PHE:HZ	1.67	0.58
1:A:130:MET:HE3	1:A:142:VAL:HG13	1.84	0.57
1:A:130:MET:HE1	1:A:195:VAL:HG11	1.87	0.54
1:A:1097:PHE:O	1:A:1100:ILE:HG12	2.11	0.51
3:C:2023:TRP:NE1	3:C:2057:MET:SD	2.86	0.49
1:A:812:TYR:CZ	2:B:241:PRO:HB3	2.50	0.46
2:B:353:HIS:NE2	5:B:502:A1BC8:S5	2.88	0.46
3:C:2023:TRP:CD1	3:C:2057:MET:SD	3.10	0.45
2:B:353:HIS:CD2	5:B:502:A1BC8:S5	3.10	0.45
2:B:350:VAL:HG22	2:B:356:VAL:HG22	2.00	0.44
1:A:392:ASN:OD1	1:A:1012:LEU:HB3	2.17	0.44
1:A:928:ARG:NH1	1:A:950:ASN:HB3	2.32	0.44
1:A:252:ILE:HD11	1:A:302:VAL:HG12	2.00	0.43
2:B:258:LYS:HE2	2:B:270:ASP:OD1	2.19	0.42
1:A:872:SER:OG	1:A:912:LEU:O	2.30	0.42
1:A:793:ILE:HG21	1:A:853:TYR:CZ	2.55	0.42
2:B:82:PRO:HG2	2:B:113:LEU:HD21	2.02	0.41
1:A:775:THR:O	1:A:777:PRO:HD3	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	799/1140 (70%)	765 (96%)	31 (4%)	3 (0%)	30	54
2	B	349/442 (79%)	331 (95%)	17 (5%)	1 (0%)	37	60
3	C	91/98 (93%)	89 (98%)	2 (2%)	0	100	100
All	All	1239/1680 (74%)	1185 (96%)	50 (4%)	4 (0%)	37	60

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	36	ASN
2	B	412	GLN
1	A	319	ASN
1	A	310	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	706/999 (71%)	687 (97%)	19 (3%)	40	64
2	B	320/398 (80%)	308 (96%)	12 (4%)	28	53
3	C	82/85 (96%)	75 (92%)	7 (8%)	8	23
All	All	1108/1482 (75%)	1070 (97%)	38 (3%)	32	57

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	THR
1	A	150	LYS
1	A	244	LYS
1	A	269	SER
1	A	282	MET
1	A	290	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	296	THR
1	A	315	THR
1	A	372	GLN
1	A	708	GLN
1	A	720	SER
1	A	823	LYS
1	A	835	MET
1	A	879	LYS
1	A	900	ARG
1	A	1006	VAL
1	A	1054	MET
1	A	1106	GLN
1	A	1135	GLU
2	B	119	THR
2	B	152	ILE
2	B	170	LEU
2	B	187	GLU
2	B	225	GLN
2	B	262	ARG
2	B	263	GLU
2	B	296	ILE
2	B	299	LEU
2	B	327	GLN
2	B	410	SER
2	B	412	GLN
3	C	2047	MET
3	C	2057	MET
3	C	2059	GLU
3	C	2064	THR
3	C	2066	LYS
3	C	2080	GLU
3	C	2084	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	105	HIS
1	A	156	ASN
1	A	255	GLN
1	A	337	ASN
1	A	708	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	790	ASN
1	A	826	ASN
1	A	907	ASN
1	A	978	GLN
1	A	1009	HIS
2	B	73	HIS
2	B	86	GLN
2	B	100	GLN
2	B	236	ASN
2	B	367	ASN
3	C	2072	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	A1BC8	B	502	-	19,22,22	0.91	1 (5%)	23,31,31	1.91	7 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	A1BC8	B	502	-	-	0/1/23/23	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	502	A1BC8	C7-C3	2.87	1.44	1.39

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	502	A1BC8	C4-C3-C7	6.03	113.35	111.69
5	B	502	A1BC8	C6-C7-C3	-2.96	110.87	111.69
5	B	502	A1BC8	C19-C13-N10	2.73	112.34	109.28
5	B	502	A1BC8	C9-N10-C13	2.41	128.29	125.62
5	B	502	A1BC8	C3-C4-S5	-2.34	111.03	112.39
5	B	502	A1BC8	C7-C8-N12	2.32	124.62	120.93
5	B	502	A1BC8	C6-C7-C8	-2.15	121.64	128.37

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	502	A1BC8	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	807/1140 (70%)	0.02	16 (1%) 64 63	41, 66, 133, 206	0
2	B	355/442 (80%)	0.37	18 (5%) 34 31	41, 76, 196, 274	0
3	C	93/98 (94%)	0.61	2 (2%) 62 60	84, 113, 164, 222	0
All	All	1255/1680 (74%)	0.16	36 (2%) 54 50	41, 71, 171, 274	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1112	LEU	3.6
1	A	770	LEU	3.4
2	B	236	ASN	3.2
2	B	128	VAL	2.8
1	A	982	ALA	2.8
2	B	144	TYR	2.7
2	B	357	HIS	2.7
2	B	97	LEU	2.7
1	A	856	GLY	2.6
2	B	297	GLN	2.5
1	A	965	PHE	2.5
1	A	920	PHE	2.4
3	C	2027	TRP	2.3
2	B	411	PRO	2.3
2	B	90	ILE	2.3
1	A	774	SER	2.3
2	B	187	GLU	2.3
1	A	1097	PHE	2.3
2	B	210	SER	2.3
2	B	102	PHE	2.3
2	B	150	PHE	2.2
1	A	49	LEU	2.2
2	B	220	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	984	THR	2.1
2	B	442	LEU	2.1
2	B	180	ALA	2.1
1	A	289	GLU	2.1
2	B	152	ILE	2.1
3	C	2113	LYS	2.1
1	A	1106	GLN	2.1
2	B	125	TYR	2.1
1	A	776	ALA	2.0
2	B	126	SER	2.0
1	A	769	LYS	2.0
1	A	370	GLN	2.0
1	A	938	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	A1BC8	B	502	20/20	0.90	0.10	55,69,91,91	0
4	ZN	B	501	1/1	0.92	0.21	251,251,251,251	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.