



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 29, 2024 – 01:28 AM EDT

PDB ID : 9WGA
Title : 2.2 ANGSTROMS RESOLUTION STRUCTURE ANALYSIS OF TWO RE-FINED N-ACETYLNEURAMINYLLACTOSE-WHEAT GERM AGGLUTININ ISOLECTIN COMPLEXES
Authors : Wright, C.S.
Deposited on : 1990-04-20
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	171	
1	B	171	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2544 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called WHEAT GERM LECTIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	171	Total	C	N	O	S	0	0	0
			1159	674	215	236	34			
1	B	171	Total	C	N	O	S	0	0	0
			1154	669	215	236	34			

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	109	Total	O	0	0
			109	109		
2	B	122	Total	O	0	0
			122	122		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: WHEAT GERM LECTIN

Chain A: 



• Molecule 1: WHEAT GERM LECTIN

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	51.34Å 73.53Å 91.54Å 90.00° 97.59° 90.00°	Depositor
Resolution (Å)	8.00 – 1.80 45.37 – 1.83	Depositor EDS
% Data completeness (in resolution range)	(Not available) (8.00-1.80) 98.4 (45.37-1.83)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	PROLSQ	Depositor
R, R_{free}	0.175 , (Not available) 0.266 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.525	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 69.4	EDS
L-test for twinning ¹	$\langle L \rangle = 0.58$, $\langle L^2 \rangle = 0.44$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2544	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.13	5/1175 (0.4%)	1.75	15/1575 (1.0%)
1	B	1.11	4/1168 (0.3%)	1.78	13/1563 (0.8%)
All	All	1.12	9/2343 (0.4%)	1.76	28/3138 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	72	GLU	CD-OE2	6.89	1.33	1.25
1	B	5	GLU	CD-OE2	6.84	1.33	1.25
1	A	5	GLU	CD-OE2	6.61	1.32	1.25
1	A	72	GLU	CD-OE1	-6.27	1.18	1.25
1	A	115	GLU	CD-OE2	6.09	1.32	1.25
1	A	72	GLU	CD-OE2	5.73	1.31	1.25
1	A	83	CYS	CB-SG	-5.65	1.72	1.81
1	B	11	GLU	CD-OE2	5.37	1.31	1.25
1	B	115	GLU	CD-OE2	5.15	1.31	1.25

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	84	ARG	NE-CZ-NH1	15.64	128.12	120.30
1	B	170	ASP	CB-CG-OD1	13.11	130.10	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	84	ARG	CD-NE-CZ	9.57	137.00	123.60
1	A	139	ARG	NE-CZ-NH2	-9.36	115.62	120.30
1	A	72	GLU	OE1-CD-OE2	-7.52	114.28	123.30
1	A	170	ASP	CB-CG-OD2	-7.40	111.64	118.30
1	B	139	ARG	CA-CB-CG	7.10	129.02	113.40
1	B	84	ARG	NE-CZ-NH2	-7.04	116.78	120.30
1	B	170	ASP	CB-CG-OD2	-6.92	112.08	118.30
1	A	85	ALA	CB-CA-C	-6.52	100.32	110.10
1	B	60	CYS	O-C-N	6.06	132.40	122.70
1	A	26	MET	CA-CB-CG	6.00	123.51	113.30
1	B	86	ASP	CB-CG-OD2	-5.96	112.94	118.30
1	B	73	TYR	CA-CB-CG	-5.95	102.10	113.40
1	A	37	ASN	CA-C-O	-5.70	108.14	120.10
1	B	123	SER	N-CA-CB	-5.56	102.16	110.50
1	A	17	CYS	O-C-N	5.54	131.56	122.70
1	B	84	ARG	CG-CD-NE	5.48	123.32	111.80
1	A	115	GLU	CG-CD-OE1	5.39	129.07	118.30
1	A	30	TYR	O-C-N	5.33	131.22	122.70
1	A	73	TYR	O-C-N	5.23	131.06	122.70
1	A	3	CYS	CA-CB-SG	5.18	123.33	114.00
1	A	115	GLU	CG-CD-OE2	-5.16	107.98	118.30
1	A	170	ASP	N-CA-CB	5.13	119.84	110.60
1	B	139	ARG	NE-CZ-NH1	5.07	122.84	120.30
1	A	99	PRO	N-CD-CG	-5.06	95.61	103.20
1	A	116	PHE	O-C-N	5.03	130.75	122.70
1	B	21	TYR	CB-CG-CD1	5.02	124.01	121.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	45	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1159	0	972	36	0
1	B	1154	0	975	43	0
2	A	109	0	0	4	0
2	B	122	0	0	10	0
All	All	2544	0	1947	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:49:GLN:NE2	1:B:79:GLN:HE22	1.60	0.99
1:A:16:LEU:HD12	1:A:26:MET:HE2	1.50	0.94
1:B:92:GLN:HE22	1:B:122:GLN:HE22	1.28	0.79
1:B:7:GLY:O	1:B:8:SER:HB2	1.84	0.78
1:B:49:GLN:HE22	1:B:79:GLN:HE22	1.32	0.76
1:A:16:LEU:HD12	1:A:26:MET:CE	2.14	0.76
1:A:49:GLN:H	1:A:49:GLN:HE21	1.34	0.75
1:A:5:GLU:HG2	2:A:274:HOH:O	1.87	0.74
1:A:49:GLN:HE22	1:A:79:GLN:HE22	1.34	0.72
1:A:49:GLN:H	1:A:49:GLN:NE2	1.88	0.71
1:A:92:GLN:HE22	1:A:122:GLN:HE22	1.39	0.70
1:B:7:GLY:O	1:B:10:MET:HB2	1.93	0.68
1:A:6:GLN:HE21	1:A:6:GLN:H	1.44	0.64
1:A:92:GLN:NE2	1:A:92:GLN:H	1.95	0.64
1:B:59:HIS:ND1	2:B:183:HOH:O	2.30	0.63
1:B:16:LEU:HD23	1:B:26:MET:CE	2.28	0.63
1:A:99:PRO:O	1:A:102:LEU:HD13	1.98	0.63
1:B:16:LEU:HD23	1:B:26:MET:HE3	1.81	0.62
1:A:20:GLN:HE22	1:A:33:LYS:NZ	1.98	0.62
1:B:2:ARG:NH2	2:B:284:HOH:O	2.32	0.61
1:B:6:GLN:NE2	1:B:6:GLN:H	1.99	0.60
1:A:45:ARG:NH1	1:A:65:GLY:O	2.34	0.60
1:A:6:GLN:HE22	1:A:36:GLN:HE22	1.50	0.58
1:B:49:GLN:HG2	1:B:65:GLY:HA3	1.85	0.58
1:B:90:GLY:C	1:B:92:GLN:H	2.08	0.56
1:A:62:SER:HB3	1:A:73:TYR:CD1	2.40	0.56
1:B:63:GLN:NE2	2:B:242:HOH:O	2.38	0.55
1:B:49:GLN:CG	1:B:65:GLY:HA3	2.37	0.55
1:A:49:GLN:NE2	1:A:79:GLN:HE22	2.05	0.54
1:A:92:GLN:H	1:A:92:GLN:HE21	1.55	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:20:GLN:HG2	2:B:265:HOH:O	2.07	0.53
1:B:63:GLN:NE2	2:B:187:HOH:O	2.37	0.53
1:A:18:CYS:HB3	1:A:36:GLN:HB2	1.91	0.53
1:B:6:GLN:HE22	1:B:36:GLN:HE22	1.56	0.53
1:A:12:CYS:HB2	2:A:218:HOH:O	2.09	0.53
1:A:45:ARG:HH12	1:A:49:GLN:HB2	1.74	0.52
1:B:47:GLY:HA2	2:B:209:HOH:O	2.10	0.51
1:B:49:GLN:NE2	1:B:49:GLN:H	2.08	0.51
1:B:96:LYS:HD3	2:B:208:HOH:O	2.10	0.51
1:B:90:GLY:C	1:B:92:GLN:N	2.64	0.50
1:A:45:ARG:HH12	1:A:49:GLN:CB	2.25	0.49
1:B:74:CYS:O	1:B:84:ARG:HB2	2.13	0.49
1:B:2:ARG:O	1:B:7:GLY:HA3	2.12	0.49
1:A:20:GLN:NE2	2:A:276:HOH:O	2.46	0.49
1:A:37:ASN:HD22	1:A:37:ASN:C	2.17	0.48
1:A:149:LYS:HA	1:A:163:GLY:O	2.14	0.47
1:B:37:ASN:ND2	2:B:213:HOH:O	2.46	0.47
1:A:20:GLN:HE22	1:A:33:LYS:HZ1	1.62	0.47
1:A:49:GLN:NE2	1:A:49:GLN:N	2.59	0.47
1:B:98:CYS:HB2	1:B:102:LEU:O	2.15	0.46
1:B:130:LYS:HB3	1:B:130:LYS:HE2	1.82	0.46
1:A:95:GLY:HA3	1:B:165:GLN:O	2.16	0.46
1:A:164:CYS:HB3	1:A:171:ALA:OXT	2.15	0.45
1:A:146:CYS:SG	1:A:168:GLY:O	2.75	0.45
1:A:130:LYS:HA	1:A:131:PRO:HD3	1.76	0.45
1:A:63:GLN:HB3	1:A:77:GLY:HA3	1.99	0.44
1:B:45:ARG:HA	1:B:65:GLY:O	2.16	0.44
1:A:16:LEU:HA	1:A:26:MET:HE2	1.98	0.44
1:B:92:GLN:H	1:B:92:GLN:HE21	1.66	0.44
1:B:83:CYS:HB2	1:B:87:ILE:HD11	1.99	0.44
1:B:33:LYS:H	1:B:33:LYS:HG3	1.59	0.44
1:B:45:ARG:HA	1:B:45:ARG:HD2	1.85	0.43
1:B:133:GLY:HA2	2:B:231:HOH:O	2.19	0.43
1:A:100:ASN:O	1:A:101:ASN:HB2	2.18	0.43
1:B:115:GLU:HG3	2:B:256:HOH:O	2.18	0.43
1:A:88:LYS:HG3	1:A:109:PHE:CE1	2.54	0.43
1:B:149:LYS:HB3	1:B:163:GLY:O	2.19	0.42
1:B:12:CYS:HB3	1:B:16:LEU:HB2	2.01	0.42
1:B:118:GLY:O	1:B:119:GLY:C	2.58	0.42
1:A:20:GLN:HE22	1:A:33:LYS:HZ3	1.65	0.42
1:B:109:PHE:HD1	1:B:109:PHE:HA	1.65	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:158:GLY:HA3	2:A:238:HOH:O	2.20	0.41
1:B:87:ILE:O	1:B:109:PHE:HA	2.21	0.41
1:A:88:LYS:HG3	1:A:109:PHE:HE1	1.86	0.41
1:B:133:GLY:O	1:B:136:ALA:N	2.40	0.41
1:B:45:ARG:HB3	1:B:50:ALA:HB2	2.02	0.41
1:B:97:LEU:HD23	1:B:97:LEU:HA	1.91	0.40
1:B:117:CYS:O	1:B:121:CYS:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	169/171 (99%)	155 (92%)	13 (8%)	1 (1%)	25	12
1	B	169/171 (99%)	158 (94%)	11 (6%)	0	100	100
All	All	338/342 (99%)	313 (93%)	24 (7%)	1 (0%)	41	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	125	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/120 (98%)	108 (92%)	10 (8%)	10	3
1	B	118/120 (98%)	102 (86%)	16 (14%)	3	1
All	All	236/240 (98%)	210 (89%)	26 (11%)	6	1

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	GLU
1	A	6	GLN
1	A	16	LEU
1	A	33	LYS
1	A	37	ASN
1	A	49	GLN
1	A	54	THR
1	A	92	GLN
1	A	97	LEU
1	A	152	SER
1	B	6	GLN
1	B	10	MET
1	B	33	LYS
1	B	37	ASN
1	B	44	LYS
1	B	49	GLN
1	B	79	GLN
1	B	87	ILE
1	B	92	GLN
1	B	93	SER
1	B	96	LYS
1	B	99	PRO
1	B	112	LEU
1	B	114	SER
1	B	139	ARG
1	B	140	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	9	ASN
1	A	20	GLN
1	A	37	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	49	GLN
1	A	63	GLN
1	A	92	GLN
1	B	6	GLN
1	B	9	ASN
1	B	14	ASN
1	B	37	ASN
1	B	49	GLN
1	B	92	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	B	1	1	7,8,9	0.82	0	9,10,12	1.46	2 (22%)
1	PCA	A	1	1	7,8,9	1.00	1 (14%)	9,10,12	1.84	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	B	1	1	-	0/0/11/13	0/1/1/1
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	PCA	O-C	2.39	1.29	1.19

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	PCA	CB-CA-C	3.86	118.02	112.70
1	B	1	PCA	O-C-CA	-3.09	116.67	124.78
1	A	1	PCA	O-C-CA	-3.05	116.78	124.78
1	B	1	PCA	CB-CA-C	2.53	116.19	112.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.