



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2024 – 09:08 am GMT

PDB ID : 4XZ3  
Title : Ca. Korarchaeum cryptofilum dinucleotide forming Acetyl-coenzyme A synthetase 1 (Se-Met derivative) in complex with coenzyme A and Mg-AMPPCP, phosphohistidine segment pointing towards nucleotide binding site  
Authors : Weisse, R.H.-J.; Scheidig, A.J.  
Deposited on : 2015-02-03  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	
1	C	464	
2	B	230	
2	D	230	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 21079 atoms, of which 10529 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acyl-CoA synthetase (NDP forming).

Mol	Chain	Residues	Atoms								ZeroOcc	AltConf	Trace
1	A	463	Total	C	H	N	O	S	Se		0	0	0
			6940	2205	3492	574	655	3	11				
1	C	461	Total	C	H	N	O	S	Se		0	0	0
			6912	2198	3476	570	654	3	11				

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	initiating methionine	UNP B1L3C9
C	1	MSE	-	initiating methionine	UNP B1L3C9

- Molecule 2 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms								ZeroOcc	AltConf	Trace
2	B	228	Total	C	H	N	O	Se			0	1	0
			3631	1145	1841	307	334	4					
2	D	213	Total	C	H	N	O	Se			0	0	0
			3225	1038	1612	276	295	4					

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MSE	-	initiating methionine	UNP B1L7P8
D	1	MSE	-	initiating methionine	UNP B1L7P8

- Molecule 3 is COENZYME A (three-letter code: COA) (formula: C<sub>21</sub>H<sub>36</sub>N<sub>7</sub>O<sub>16</sub>P<sub>3</sub>S).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	P	S	0	0
			80	21	32	7	16	3	1		
3	C	1	Total	C	H	N	O	P	S	0	0
			80	21	32	7	16	3	1		

- Molecule 4 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (three-letter code: ACP) (formula: C<sub>11</sub>H<sub>18</sub>N<sub>5</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	P		0	0
			31	6	9	1	12	3			

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	H	N	O	P	
			44	11	13	5	12	3	
4	C	1	Total	C	H	N	O	P	
			31	6	9	1	12	3	
4	D	1	Total	C	H	N	O	P	
			44	11	13	5	12	3	

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg		
			1	1	0	0
5	B	1	Total	Mg		
			1	1	0	0
5	C	2	Total	Mg		
			2	2	0	0
5	D	1	Total	Mg		
			1	1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	23	Total	O		
			23	23	0	0
6	B	11	Total	O		
			11	11	0	0
6	C	21	Total	O		
			21	21	0	0
6	D	1	Total	O		
			1	1	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acyl-CoA synthetase (NDP forming)




- Molecule 1: Acyl-CoA synthetase (NDP forming)



- Molecule 2: Uncharacterized protein



- Molecule 2: Uncharacterized protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.17Å 111.88Å 127.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.84 – 2.40 48.84 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.84-2.40) 99.8 (48.84-2.40)	Depositor EDS
$R_{merge}$	0.27	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.19 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, $R_{free}$	0.197 , 0.244 0.203 , 0.247	Depositor DCC
$R_{free}$ test set	2878 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.4	Xtriage
Anisotropy	0.087	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 35.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	21079	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACP, MG, COA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.25	0/3505	0.40	0/4743
1	C	0.25	0/3493	0.40	0/4729
2	B	0.25	0/1817	0.44	0/2445
2	D	0.25	0/1634	0.44	0/2202
All	All	0.25	0/10449	0.41	0/14119

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3448	3492	3494	12	0
1	C	3436	3476	3478	8	0
2	B	1790	1841	1843	11	0
2	D	1613	1612	1614	11	0
3	A	48	32	32	1	0
3	C	48	32	32	0	0
4	A	22	9	10	1	0
4	B	31	13	14	1	0
4	C	22	9	10	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	31	13	14	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	2	0	0	0	0
5	D	1	0	0	0	0
6	A	23	0	0	0	0
6	B	11	0	0	0	0
6	C	21	0	0	0	0
6	D	1	0	0	0	0
All	All	10550	10529	10541	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:GLU:OE2	1:A:116:ARG:NH1	2.36	0.58
2:B:173:LEU:HG	2:D:145:LEU:HD11	1.88	0.55
1:A:187:LYS:NZ	1:A:192:ASP:OD1	2.31	0.51
1:C:359:ASP:OD2	1:C:359:ASP:N	2.32	0.50
2:B:51:ARG:NH2	2:B:82:GLU:OE1	2.45	0.49
1:A:343:PHE:O	4:A:502:ACP:O3'	2.27	0.48
2:B:3:SER:OG	2:B:5:ASP:OD1	2.25	0.48
2:B:124:LEU:HB3	2:B:208:MSE:HB3	1.96	0.47
2:B:177:PHE:HZ	2:D:144:GLU:HG2	1.81	0.46
2:B:43:GLU:OE2	2:B:90:ARG:NH2	2.49	0.46
1:A:237:ILE:HD11	1:A:290:LEU:HD12	1.96	0.45
2:B:24:LYS:NZ	2:B:223:VAL:O	2.43	0.45
1:C:237:ILE:HD11	1:C:290:LEU:HD12	1.98	0.45
1:A:328:PRO:HD3	1:A:348:ASN:HB2	1.99	0.44
2:D:126:ARG:NH2	2:D:202:ASN:O	2.51	0.43
1:A:66:ASP:OD1	1:A:67:LYS:N	2.51	0.43
2:D:67:VAL:HG13	2:D:68:HIS:N	2.34	0.43
1:A:244:THR:HG21	1:C:319:ALA:HB3	2.01	0.43
1:A:341:PRO:HD3	1:A:360:TRP:CH2	2.53	0.43
1:C:200:ASP:O	1:C:233:ARG:NH2	2.52	0.43
2:B:138:LEU:HD21	2:B:173:LEU:HD12	2.01	0.43
2:D:69:LYS:HB3	2:D:75:VAL:HG23	2.01	0.43
1:C:136:ALA:N	1:C:137:PRO:CD	2.82	0.42
1:A:215:ILE:CD1	1:A:220:GLY:HA2	2.50	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:226:ARG:NH2	4:B:300:ACP:O3G	2.52	0.42
1:C:215:ILE:CD1	1:C:220:GLY:HA2	2.49	0.42
2:D:228:ILE:HD12	2:D:228:ILE:N	2.35	0.42
1:A:161:GLY:HA3	3:A:501:COA:S1P	2.59	0.42
2:D:136:PHE:CG	2:D:137:GLY:N	2.88	0.42
2:D:124:LEU:HB3	2:D:208:MSE:HB3	2.02	0.41
2:B:145:LEU:HD21	2:D:170:ALA:HA	2.02	0.41
2:D:37:ILE:HG22	2:D:38:ARG:N	2.35	0.41
2:D:67:VAL:HG13	2:D:68:HIS:H	1.85	0.41
1:C:246:VAL:HG22	1:C:247:GLY:N	2.36	0.41
1:A:136:ALA:N	1:A:137:PRO:CD	2.85	0.40
1:C:290:LEU:HD23	1:C:447:LEU:HD23	2.03	0.40
2:B:89:TYR:CE1	2:B:108:ILE:HG13	2.56	0.40
1:A:290:LEU:HD23	1:A:447:LEU:HD23	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	461/464 (99%)	446 (97%)	14 (3%)	1 (0%)	44	59
1	C	459/464 (99%)	441 (96%)	17 (4%)	1 (0%)	44	59
2	B	227/230 (99%)	220 (97%)	7 (3%)	0	100	100
2	D	207/230 (90%)	201 (97%)	6 (3%)	0	100	100
All	All	1354/1388 (98%)	1308 (97%)	44 (3%)	2 (0%)	48	65

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	348	ASN

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	348	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	366/358 (102%)	360 (98%)	6 (2%)	58	76
1	C	365/358 (102%)	359 (98%)	6 (2%)	58	76
2	B	194/192 (101%)	192 (99%)	2 (1%)	73	86
2	D	165/192 (86%)	164 (99%)	1 (1%)	84	92
All	All	1090/1100 (99%)	1075 (99%)	15 (1%)	62	79

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	192	ASP
1	A	196	LEU
1	A	222	MSE
1	A	257	SER
1	A	258	ILE
1	A	355	MSE
2	B	146	PHE
2	B	169	LYS
1	C	6	ARG
1	C	112	GLU
1	C	276	MSE
1	C	355	MSE
1	C	359	ASP
1	C	402	GLU
2	D	80	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	COA	A	501	-	41,50,50	0.54	0	52,75,75	0.59	1 (1%)
3	COA	C	504	-	41,50,50	0.49	0	52,75,75	0.59	1 (1%)
4	ACP	B	300	5	27,33,33	0.91	2 (7%)	32,52,52	1.14	4 (12%)
4	ACP	C	501	5	18,22,33	1.04	3 (16%)	25,35,52	1.14	3 (12%)
4	ACP	A	502	5	18,22,33	0.86	2 (11%)	25,35,52	1.16	3 (12%)
4	ACP	D	300	5	27,33,33	1.01	2 (7%)	32,52,52	1.09	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	COA	A	501	-	-	6/44/64/64	0/3/3/3
3	COA	C	504	-	-	2/44/64/64	0/3/3/3

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ACP	B	300	5	-	1/15/38/38	0/3/3/3
4	ACP	C	501	5	-	7/15/34/38	0/1/1/3
4	ACP	A	502	5	-	0/15/34/38	0/1/1/3
4	ACP	D	300	5	-	5/15/38/38	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	501	ACP	PG-O1G	2.30	1.55	1.50
4	B	300	ACP	PG-O1G	2.25	1.54	1.50
4	D	300	ACP	PG-O1G	2.25	1.54	1.50
4	A	502	ACP	PG-O1G	2.23	1.54	1.50
4	D	300	ACP	PG-O2G	-2.20	1.49	1.54
4	B	300	ACP	PG-O3G	-2.15	1.50	1.54
4	C	501	ACP	PG-O3G	-2.15	1.50	1.54
4	A	502	ACP	PG-O2G	-2.15	1.50	1.54
4	C	501	ACP	PB-O1B	2.00	1.56	1.51

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	502	ACP	O1B-PB-C3B	3.67	118.77	109.07
4	B	300	ACP	O1B-PB-C3B	3.64	118.71	109.07
4	C	501	ACP	O2B-PB-C3B	3.24	119.84	106.58
4	D	300	ACP	O2B-PB-C3B	3.09	119.20	106.58
4	B	300	ACP	O1G-PG-C3B	-2.49	105.87	111.24
4	C	501	ACP	O3G-PG-C3B	2.49	112.44	106.40
4	C	501	ACP	O1G-PG-C3B	-2.42	106.02	111.24
4	B	300	ACP	O3G-PG-C3B	2.42	112.26	106.40
4	A	502	ACP	O1G-PG-C3B	-2.40	106.07	111.24
4	D	300	ACP	O1G-PG-C3B	-2.39	106.10	111.24
4	D	300	ACP	O2G-PG-C3B	2.35	112.11	106.40
4	A	502	ACP	O2G-PG-C3B	2.34	112.07	106.40
3	C	504	COA	C5A-C6A-N6A	2.29	123.83	120.35
4	D	300	ACP	C5-C6-N6	2.23	123.74	120.35
3	A	501	COA	C5A-C6A-N6A	2.22	123.73	120.35
4	B	300	ACP	C5-C6-N6	2.21	123.71	120.35

There are no chirality outliers.

All (21) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	C	501	ACP	C5'-O5'-PA-O1A
4	C	501	ACP	C5'-O5'-PA-O2A
4	D	300	ACP	PG-C3B-PB-O1B
4	D	300	ACP	PG-C3B-PB-O2B
4	D	300	ACP	C5'-O5'-PA-O1A
4	D	300	ACP	C5'-O5'-PA-O2A
4	D	300	ACP	C5'-O5'-PA-O3A
3	C	504	COA	C6P-C7P-N8P-C9P
3	A	501	COA	C3B-C4B-C5B-O5B
3	A	501	COA	O4B-C4B-C5B-O5B
4	C	501	ACP	O4'-C4'-C5'-O5'
4	C	501	ACP	C3'-C4'-C5'-O5'
3	C	504	COA	P2A-O3A-P1A-O1A
3	A	501	COA	S1P-C2P-C3P-N4P
4	C	501	ACP	PB-O3A-PA-O5'
3	A	501	COA	C3B-O3B-P3B-O7A
3	A	501	COA	C3B-O3B-P3B-O9A
4	C	501	ACP	PB-C3B-PG-O3G
3	A	501	COA	C4B-C5B-O5B-P1A
4	C	501	ACP	C5'-O5'-PA-O3A
4	B	300	ACP	C5'-O5'-PA-O1A



There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	COA	1	0
4	B	300	ACP	1	0
4	A	502	ACP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	452/464 (97%)	-0.00	9 (1%) 64 61	32, 52, 88, 119	0
1	C	450/464 (96%)	-0.06	9 (2%) 64 61	30, 52, 87, 138	0
2	B	224/230 (97%)	0.29	6 (2%) 56 53	36, 69, 105, 128	1 (0%)
2	D	209/230 (90%)	1.12	39 (18%) 4 4	41, 87, 145, 180	0
All	All	1335/1388 (96%)	0.20	63 (4%) 37 34	30, 58, 114, 180	1 (0%)

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	108	ILE	4.7
2	D	45	GLU	4.5
2	D	95	ASN	4.4
2	D	64	PRO	4.3
2	D	109	LEU	4.0
2	D	49	PHE	3.5
2	D	142	PHE	3.5
2	D	73	GLY	3.5
2	D	62	VAL	3.4
2	D	92	ILE	3.4
2	D	89	TYR	3.4
2	D	39	LEU	3.4
2	D	61	ILE	3.3
2	D	47	VAL	3.2
2	D	146	PHE	3.2
2	D	50	SER	3.1
1	C	463	VAL	3.1
1	C	248	ALA	3.0
2	D	44	GLU	3.0
2	D	46	ALA	2.9
1	C	246	VAL	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	D	67	VAL	2.9
1	C	255	THR	2.9
2	D	72	VAL	2.8
2	B	177	PHE	2.8
2	D	58	VAL	2.8
2	D	145	LEU	2.8
2	D	83	GLU	2.7
1	A	81	LEU	2.6
1	A	246	VAL	2.6
2	D	177	PHE	2.6
2	B	146	PHE	2.5
2	D	118	VAL	2.5
2	D	60	LYS	2.5
1	C	257	SER	2.5
2	D	63	SER	2.5
2	D	74	GLY	2.5
2	D	141	VAL	2.5
2	D	143	VAL	2.4
2	B	101	PRO	2.4
1	A	85	ASP	2.4
2	B	67	VAL	2.4
2	D	65	GLN	2.4
2	D	179	GLY	2.4
2	D	59	LEU	2.4
1	A	58	TYR	2.4
2	D	52	GLU	2.3
2	B	179	GLY	2.3
2	D	93	ILE	2.3
1	A	2	ASN	2.2
1	A	464	ARG	2.2
2	B	142	PHE	2.2
2	D	229	LEU	2.2
1	C	259	ALA	2.2
2	D	125	LEU	2.1
1	C	252	ALA	2.1
2	D	55	PHE	2.1
1	A	247	GLY	2.1
1	C	258	ILE	2.1
1	A	248	ALA	2.1
1	C	250	ALA	2.1
2	D	110	VAL	2.0
1	A	78	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	ACP	D	300	31/31	0.75	0.12	91,134,166,177	0
4	ACP	A	502	22/31	0.83	0.12	63,91,115,123	0
5	MG	C	503	1/1	0.84	0.32	59,59,59,59	0
5	MG	C	502	1/1	0.86	0.27	62,62,62,62	0
5	MG	D	301	1/1	0.87	0.10	93,93,93,93	0
3	COA	A	501	48/48	0.91	0.10	52,69,81,93	0
3	COA	C	504	48/48	0.92	0.09	46,64,79,84	0
4	ACP	C	501	22/31	0.92	0.09	53,82,107,115	0
5	MG	A	503	1/1	0.93	0.17	57,57,57,57	0
5	MG	B	301	1/1	0.94	0.11	57,57,57,57	0
4	ACP	B	300	31/31	0.94	0.07	51,69,85,90	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around ACP D 300:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around ACP A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around COA A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around COA C 504:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around ACP C 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ACP B 300:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.