



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 03:28 PM EDT

PDB ID : 6ZAT  
Title : Nitrite-bound copper nitrite reductase from Bradyrhizobium sp. ORS 375 (two-domain) at 1.0 Å resolution (unrestrained full matrix refinement by SHELX)  
Authors : Rose, S.L.; Antonyuk, S.V.; Sasaki, D.; Yamashita, K.; Hirata, K.; Ueno, G.; Ago, H.; Eady, R.R.; Tosha, T.; Yamamoto, M.; Hasnain, S.S.  
Deposited on : 2020-06-05  
Resolution : 1.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 1.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1050 (1.06-0.94)
Clashscore	141614	1117 (1.06-0.94)
Ramachandran outliers	138981	1043 (1.06-0.94)
Sidechain outliers	138945	1045 (1.06-0.94)
RSRZ outliers	127900	1023 (1.06-0.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	504[A]	-	X	-	-

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3627 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Copper-containing nitrite reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	345	Total	C	N	O	S	0	27	0
			2776	1772	476	514	14			


There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP H0SLX7
A	342	GLU	-	expression tag	UNP H0SLX7
A	343	ASN	-	expression tag	UNP H0SLX7
A	344	LEU	-	expression tag	UNP H0SLX7
A	345	TYR	-	expression tag	UNP H0SLX7
A	346	PHE	-	expression tag	UNP H0SLX7
A	347	GLN	-	expression tag	UNP H0SLX7

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cu	0	1
			3	3		

- Molecule 3 is NITRITE ION (three-letter code: NO2) (formula: NO<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	N	O	0	1
			6	2	4		
3	A	1	Total	N	O	0	1
			3	1	2		
3	A	1	Total	N	O	0	1
			6	2	4		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	1
			6	3	3		


- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	1
			5	4	1		
5	A	1	Total	O	S	0	1
			5	4	1		
5	A	1	Total	O	S	0	1
			5	4	1		
5	A	1	Total	O	S	0	1
			5	4	1		
5	A	1	Total	O	S	0	1
			5	4	1		
5	A	1	Total	O	S	0	1
			5	4	1		
5	A	1	Total	O	S	0	1
			5	4	1		
5	A	1	Total	O	S	0	1
			5	4	1		

- Molecule 6 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID

(three-letter code: EPE) (formula:  $C_8H_{18}N_2O_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	1
			15	8	2	4	1		
6	A	1	Total	C	N	O	S	0	1
			15	8	2	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	700	Total	O	0	102
			737	737		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Copper-containing nitrite reductase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.79Å 106.79Å 106.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.00 53.39 – 1.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.00-1.00) 99.9 (53.39-1.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.26 (at 1.00Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.126 , 0.143 0.126 , 0.126	Depositor DCC
$R_{free}$ test set	10771 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	8.5	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 79.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.023 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	3627	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EPE, GOL, NO2, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.72	2/2922 (0.1%)	1.32	30/3969 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	342	GLU	CD-OE2	7.37	1.33	1.25
1	A	117	ARG	CD-NE	-5.43	1.37	1.46

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	169[A]	ARG	NE-CZ-NH2	-18.27	111.16	120.30
1	A	169[B]	ARG	NE-CZ-NH2	-18.27	111.16	120.30
1	A	117	ARG	NE-CZ-NH2	15.62	128.11	120.30
1	A	117	ARG	NE-CZ-NH1	-13.33	113.63	120.30
1	A	154	ARG	CD-NE-CZ	10.82	138.75	123.60
1	A	169[A]	ARG	NE-CZ-NH1	10.64	125.62	120.30
1	A	169[B]	ARG	NE-CZ-NH1	10.64	125.62	120.30
1	A	182	ARG	NE-CZ-NH1	10.28	125.44	120.30
1	A	8	ARG	NE-CZ-NH1	10.17	125.38	120.30
1	A	8	ARG	NE-CZ-NH2	-9.25	115.68	120.30
1	A	218[A]	LYS	CA-CB-CG	8.09	131.19	113.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218[B]	LYS	CA-CB-CG	8.09	131.19	113.40
1	A	9	GLN	CA-CB-CG	8.08	131.19	113.40
1	A	154	ARG	NE-CZ-NH1	-7.85	116.38	120.30
1	A	182	ARG	NH1-CZ-NH2	-7.32	111.34	119.40
1	A	154	ARG	CG-CD-NE	-7.25	96.56	111.80
1	A	346	PHE	CB-CG-CD2	-6.74	116.08	120.80
1	A	9	GLN	CB-CG-CD	6.54	128.61	111.60
1	A	53[A]	PHE	N-CA-CB	6.31	121.96	110.60
1	A	53[B]	PHE	N-CA-CB	6.31	121.96	110.60
1	A	346	PHE	CD1-CG-CD2	6.16	126.31	118.30
1	A	86	THR	CA-CB-CG2	-5.91	104.13	112.40
1	A	10	ARG	CD-NE-CZ	5.88	131.84	123.60
1	A	182	ARG	NE-CZ-NH2	5.79	123.19	120.30
1	A	245	ARG	NE-CZ-NH2	5.56	123.08	120.30
1	A	346	PHE	CG-CD1-CE1	-5.55	114.70	120.80
1	A	342	GLU	CG-CD-OE2	-5.52	107.26	118.30
1	A	206	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	A	87[B]	MET	CG-SD-CE	5.31	108.70	100.20
1	A	87[C]	MET	CG-SD-CE	5.31	108.70	100.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	299	VAL	Peptide
1	A	301	HIS	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2776	0	2695	16	0
2	A	3	0	0	0	0
3	A	15	0	0	1	0
4	A	6	0	7	2	0
5	A	60	0	0	0	0
6	A	30	0	34	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	737	0	0	3	0
All	All	3627	0	2736	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53[B]:PHE:HE1	7:A:777[B]:HOH:O	1.54	0.88
1:A:295:ILE:HD12	1:A:316:LYS:HG2	1.71	0.73
6:A:520[A]:EPE:H82	7:A:1068[A]:HOH:O	1.97	0.64
1:A:224:GLY:H	4:A:504[A]:GOL:H31	1.65	0.60
1:A:295:ILE:CD1	1:A:316:LYS:HG2	2.34	0.57
1:A:53[B]:PHE:CE1	7:A:777[B]:HOH:O	2.41	0.50
1:A:235:ASN:OD1	1:A:287[B]:LEU:HG	2.13	0.48
1:A:87[C]:MET:HG2	1:A:140:HIS:CE1	2.49	0.48
1:A:224:GLY:H	4:A:504[A]:GOL:C3	2.26	0.47
1:A:39:VAL:HG11	1:A:89:HIS:CD2	2.51	0.45
1:A:184:GLU:OE1	1:A:184:GLU:HA	2.15	0.45
1:A:43[A]:LYS:HE2	3:A:517[A]:NO2:N	2.32	0.45
1:A:5:LYS:HD2	1:A:5:LYS:HA	1.37	0.44
1:A:252:ILE:HG13	1:A:297[A]:ALA:O	2.19	0.42
1:A:43[A]:LYS:HE3	1:A:54:GLN:NE2	2.34	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/347 (106%)	361 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	302/283 (107%)	294 (97%)	8 (3%)	46 13

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASP
1	A	5	LYS
1	A	9	GLN
1	A	10	ARG
1	A	53[A]	PHE
1	A	53[B]	PHE
1	A	248	ARG
1	A	330[A]	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	229	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 3 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	SO4	A	515[C]	-	4,4,4	0.65	0	6,6,6	0.12	0
5	SO4	A	505[A]	-	4,4,4	0.81	0	6,6,6	0.32	0
5	SO4	A	512[A]	-	4,4,4	0.57	0	6,6,6	0.18	0
6	EPE	A	519[A]	-	15,15,15	1.31	2 (13%)	19,20,20	1.61	2 (10%)
4	GOL	A	504[A]	-	5,5,5	1.74	2 (40%)	5,5,5	2.97	3 (60%)
3	NO2	A	518[B]	-	1,2,2	0.68	0	0,1,1	-	-
3	NO2	A	503[B]	2	1,2,2	1.53	0	0,1,1	-	-
5	SO4	A	510[A]	-	4,4,4	0.69	0	6,6,6	0.18	0
5	SO4	A	509[A]	-	4,4,4	0.68	0	6,6,6	0.19	0
3	NO2	A	517[A]	-	1,2,2	0.18	0	0,1,1	-	-
3	NO2	A	518[A]	-	1,2,2	1.26	0	0,1,1	-	-
3	NO2	A	503[A]	2	1,2,2	0.58	0	0,1,1	-	-
5	SO4	A	511[A]	-	4,4,4	0.53	0	6,6,6	0.28	0
5	SO4	A	508[A]	-	4,4,4	0.58	0	6,6,6	0.15	0
5	SO4	A	506[A]	-	4,4,4	0.61	0	6,6,6	0.09	0
5	SO4	A	516[C]	-	4,4,4	0.62	0	6,6,6	0.08	0
6	EPE	A	520[A]	-	15,15,15	1.27	3 (20%)	19,20,20	2.36	6 (31%)
5	SO4	A	513[A]	-	4,4,4	0.67	0	6,6,6	0.28	0
5	SO4	A	507[A]	-	4,4,4	0.64	0	6,6,6	0.07	0
5	SO4	A	514[A]	-	4,4,4	0.59	0	6,6,6	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EPE	A	520[A]	-	-	2/9/19/19	0/1/1/1
6	EPE	A	519[A]	-	-	0/9/19/19	0/1/1/1
4	GOL	A	504[A]	-	-	3/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	520[A]	EPE	C9-N1	-2.89	1.40	1.47
4	A	504[A]	GOL	O3-C3	-2.87	1.30	1.42
6	A	519[A]	EPE	O3S-S	-2.58	1.37	1.47
6	A	520[A]	EPE	O2S-S	2.58	1.52	1.45
6	A	519[A]	EPE	C10-S	-2.48	1.74	1.77
4	A	504[A]	GOL	C1-C2	2.10	1.59	1.51
6	A	520[A]	EPE	O3S-S	-2.02	1.39	1.47

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	520[A]	EPE	O2S-S-C10	-6.19	97.37	106.73
4	A	504[A]	GOL	O3-C3-C2	5.18	133.68	110.38
6	A	519[A]	EPE	O1S-S-C10	-4.62	99.75	106.73
6	A	520[A]	EPE	C9-N1-C2	4.30	122.70	111.24
4	A	504[A]	GOL	C3-C2-C1	-3.44	99.19	111.80
6	A	520[A]	EPE	C10-C9-N1	3.27	124.75	112.36
6	A	520[A]	EPE	C9-N1-C6	3.08	119.45	111.24
6	A	520[A]	EPE	C7-N4-C3	2.84	118.80	111.24
6	A	519[A]	EPE	C5-C6-N1	2.24	115.16	110.65
6	A	520[A]	EPE	O3S-S-C10	2.20	110.30	106.00
4	A	504[A]	GOL	O2-C2-C3	-2.17	100.21	109.18

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	520[A]	EPE	N4-C7-C8-O8
4	A	504[A]	GOL	O1-C1-C2-C3
4	A	504[A]	GOL	O1-C1-C2-O2
4	A	504[A]	GOL	O2-C2-C3-O3
6	A	520[A]	EPE	C8-C7-N4-C5

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	504[A]	GOL	2	0
3	A	517[A]	NO2	1	0
6	A	520[A]	EPE	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	345/347 (99%)	-0.15	7 (2%) 65 55	6, 9, 22, 63	20 (5%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	337	THR	4.1
1	A	338	GLY	3.9
1	A	339	SER	3.4
1	A	3	ASP	2.7
1	A	345	TYR	2.4
1	A	4	LEU	2.3
1	A	346	PHE	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	SO4	A	516[C]	5/5	0.70	0.18	61,61,63,65	5

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	SO4	A	514[A]	5/5	0.82	0.25	40,63,65,69	5
4	GOL	A	504[A]	6/6	0.82	0.71	10,17,20,37	6
5	SO4	A	513[A]	5/5	0.88	0.28	11,13,14,18	5
5	SO4	A	509[A]	5/5	0.91	0.12	23,28,39,46	5
5	SO4	A	508[A]	5/5	0.91	0.14	11,14,21,27	5
5	SO4	A	515[C]	5/5	0.92	0.14	32,34,48,49	5
5	SO4	A	512[A]	5/5	0.92	0.14	12,15,20,22	5
6	EPE	A	519[A]	15/15	0.92	0.13	12,13,19,20	15
5	SO4	A	506[A]	5/5	0.95	0.18	28,33,52,60	5
5	SO4	A	510[A]	5/5	0.95	0.11	31,36,46,55	5
5	SO4	A	507[A]	5/5	0.96	0.14	35,39,57,60	5
3	NO2	A	517[A]	3/3	0.96	0.12	15,15,26,28	0
3	NO2	A	503[A]	3/3	0.96	0.10	12,12,12,13	3
5	SO4	A	505[A]	5/5	0.96	0.09	30,34,46,48	5
3	NO2	A	503[B]	3/3	0.96	0.10	12,12,13,15	3
3	NO2	A	518[B]	3/3	0.97	0.10	10,10,16,26	3
3	NO2	A	518[A]	3/3	0.97	0.10	14,14,18,24	3
5	SO4	A	511[A]	5/5	0.97	0.10	17,19,23,29	5
6	EPE	A	520[A]	15/15	0.99	0.08	6,8,26,32	15
2	CU	A	502	1/1	1.00	0.04	7,7,7,7	0
2	CU	A	501[A]	1/1	1.00	0.04	7,7,7,7	1
2	CU	A	501[B]	1/1	1.00	0.04	8,8,8,8	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around CU A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around CU A 501 (A):**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around CU A 501 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**6.5 Other polymers** ⓘ

There are no such residues in this entry.